

University of Windsor

Scholarship at UWindsor

Electronic Theses and Dissertations

Theses, Dissertations, and Major Papers

10-5-2017

Acoustic signalling in Savannah Sparrows, *Passerculus sandwichensis*: Diel and seasonal variation, male-male vocal interactions, and responses to playback

Ines Moran
University of Windsor

Follow this and additional works at: <https://scholar.uwindsor.ca/etd>

Recommended Citation

Moran, Ines, "Acoustic signalling in Savannah Sparrows, *Passerculus sandwichensis*: Diel and seasonal variation, male-male vocal interactions, and responses to playback" (2017). *Electronic Theses and Dissertations*. 7281.

<https://scholar.uwindsor.ca/etd/7281>

This online database contains the full-text of PhD dissertations and Masters' theses of University of Windsor students from 1954 forward. These documents are made available for personal study and research purposes only, in accordance with the Canadian Copyright Act and the Creative Commons license—CC BY-NC-ND (Attribution, Non-Commercial, No Derivative Works). Under this license, works must always be attributed to the copyright holder (original author), cannot be used for any commercial purposes, and may not be altered. Any other use would require the permission of the copyright holder. Students may inquire about withdrawing their dissertation and/or thesis from this database. For additional inquiries, please contact the repository administrator via email (scholarship@uwindsor.ca) or by telephone at 519-253-3000ext. 3208.

**Acoustic signalling in Savannah Sparrows, *Passerculus sandwichensis*:
Diel and seasonal variation, male-male vocal interactions,
and responses to playback**

By
Ines Géraldine Moran

A Thesis
Submitted to the Faculty of Graduate Studies
through the Department of Biological Sciences in Partial Fulfillment of the
Requirements for the Degree of Master of Science
at the University of Windsor

Windsor, Ontario, Canada
2017

© 2017 Ines G. Moran

**Acoustic signalling in Savannah Sparrows, *Passerculus sandwichensis*:
Diel and seasonal variation, male-male vocal interactions,
and responses to playback**

by

Ines Moran

APPROVED BY:

C. J. Miller,
Department of Psychology

S. M. Doucet,
Department of Biological Sciences

D. R. Norris, Co-Advisor
Department of Integrative Biology, University of Guelph

D. J. Mennill, Advisor
Department of Biological Sciences

August 28, 2017

Declaration of co-authorship

I hereby declare that this thesis incorporates material that is the result of joint research, as follows: I am the sole the author of chapter 1 and 4 and the principal author of chapters 2 and 3. Both chapters 2 and 3 were conducted under the supervision of Daniel Mennill (University of Windsor) and Ryan Norris (University of Guelph), and the guidance of two additional co-authors, Stéphanie Doucet (University of Windsor) and Amy Newman (University of Guelph), who contributed input on experimental design, analyses, and writing as well as providing funding and logistical support for field research. In all cases, the key ideas, data analysis, interpretation, and writing were performed by myself, and my co-authors provided assistance with experimental design, refinement of ideas, and editing the manuscripts.

I am aware of the University of Windsor Senate Policy on Authorship and I certify that I have properly acknowledged the contribution of other researchers to my thesis, and have obtained written permission from each of the co-author(s) to include the above material(s) in my thesis.

I certify that, with the above qualification, this thesis, and the research to which it refers, is the product of my own work.

I declare that, to the best of my knowledge, my thesis does not infringe upon anyone's copyright nor violate any proprietary rights and that any ideas, techniques, quotations, or any other material from the work of other people included in my thesis,

published or otherwise, are fully acknowledged in accordance with the standard referencing practices.

I declare that this is a true copy of my thesis, including any final revisions, as approved by my thesis committee and the Graduate Studies office, and that this thesis has not been submitted for a higher degree to any other university or institution.

Abstract

In this thesis, I examine the function of acoustic signals in Savannah Sparrows, *Passerculus sandwichensis*, by analyzing temporal variation in vocal activity, and by conducting a playback experiment. In my first data chapter, I analyze long-term acoustic recordings to study diel and seasonal variation in vocal activity of Savannah Sparrows. I show that singing activity of male Savannah Sparrows varies with time of day, time of year, and breeding stage. Males exhibit the highest level of song output in May, upon arrival on the breeding grounds, and the lowest level in August, before departure from the breeding grounds. Song output peaks in the early morning, consistent with dawn chorus behaviour, but this pattern is common only prior to pairing; after pairing the dawn chorus is reduced and male song output peaks in the evening, consistent with dusk chorus behaviour. These patterns suggest that dawn choruses serve a territorial function whereas dusk choruses serve a female-related function in Savannah Sparrows. In my second data chapter, I present the results of a playback experiment designed to test whether Savannah Sparrows signal their intention to attack a rival male. I simulated an intruder using song playback and a taxidermic model, and explored which behaviors were associated with physical attack. Savannah Sparrows produce soft songs and chip calls at significantly higher levels before attacking rivals, whereas three other measured behaviors (aggressive calls, wing waving, and passes over the model) do not predict attack. My research advances the field of animal communication and provides a foundation for future research on signal function and social interactions involving vocal signals.

Acknowledgments

I thank my supervisor Dan Mennill and my co-supervisor Ryan Norris for their help, support, and encouragement throughout my Master's degree. I thank Dan and Ryan for entrusting me to conduct field research on Kent Island in April and May of each year of my thesis; thanks to both of them, I had the privilege to work in a remarkable part of the world with a unique biological system and incredible birds.

I am extremely grateful to Dan who gave me the opportunity to explore the world of animal communication in his lab during my Master's degree. This unique opportunity brought me closer to my goal of becoming a field biologist and enabled me to gain many important skills as a biologist, scientist, and writer. I thank Dan for his generous financial support that funded my research and my participation in three conferences during my Master's degree. Finally, I would like to thank Dan for his enthusiasm, his guidance, his encouragement, and friendship over the last two years.

I thank my committee members, Ryan Norris, Stéphanie Doucet, and Carlin Miller for their enthusiasm and feedback on my thesis. I would also like to thank my collaborators, Ryan Norris, Amy Newman, Stéphanie Doucet, and Brad Woodworth for monitoring the population of Savannah Sparrows on Kent Island as well as their invaluable contribution to my research.

I would like to thank the Natural Sciences and Engineering Research Council of Canada (NSERC), the Canada Foundation for Innovation, and the University of Windsor

for supporting my research through grants to my supervisor. Further, I thank NSERC for their ongoing contributions to the research of my collaborators Ryan Norris, Amy Newman, and Stéphanie Doucet, which also helped my research indirectly. Additionally, I would like to thank the University of Windsor, Nature Vancouver, the American Ornithologists' Union, the Animal Behavior Society for their financial support through the grants and scholarships they have provided to me. I would also like to thank the Beaty Biodiversity Museum for their generous donation of taxidermic models for my second data chapter.

My research field seasons would not have been possible without the logistical support of Bowdoin Scientific Station on Kent Island. In particular, I thank Mark Murray, Russel Ingalls, Damon Gannon, Janet Gannon, Don Dearborn, Dre Gager, Ed Minot, and Midge Minot. In addition, I am thankful to Jim Leslie at the Marathon Inn.

For their valuable contribution to field research, I thank Katrina Lukianchuk, Helmi Hess, Kristina Hick, Rachel Hasson, Katherine McGuire, and Ian Thomas. For their valuable contribution to the ongoing field research on the Savannah Sparrow population on Kent Island, I thank Ryan Norris, Amy Newman, Stéphanie Doucet, Dan Mennill, and Brad Woodworth.

I would like to thank everyone in the Mennill Sound Analysis Lab for the great times spent inside and outside the lab, either birding, or socializing. I particularly thank Brendan Graham, Alana Demko, Zach Kahn, Rachel Hasson, Katrina Switzer, Lincoln Savi,

Katherine McGuire, Ian Thomas, Christopher Moser-Purdy, Kristina Hick, Blaine Landsborough, Justin Johnson, Kelly Schoenberger, and Vivian Hui.

I would like to thank the Department of Biological Sciences' office staff, Dennis Higgs, Nancy Barkley, Martha Hiuser, Judy Janzen, and Chris Côté, for their excellent service and their friendliness throughout my degree.

Finally, I would like to give a special thanks to my friends and family whose sheltering support has been very important throughout my degree. In particular, I would like to thank Kiruthika Rathanaswami, Catherine Georges, Jai Govinda, Mark Murray, Shelagh Dodd, Sue Wong, Ildiko Szabo, David Turner, Kelly Sekhon, Alice Beauvisage, Brendan Graham, Alana Demko, Meghan Mickle, Mellissa Kaitlyn, Chantal Dings, Helmi Hess, Deepika Santa Kumar, Katherine McGuire, and Ian Thomas for their generosity with their time, kind words of friendship, and encouragement. Additionally, I would also like to acknowledge a special friend, Shelvin Prakasan, who supported me at the start of this special journey. Finally, I would like to thank the most inspiring and supportive person in my life, my mother, Christine Moran. She supported me throughout my degree and encouraged me with her words of wisdom to pursue my dreams. Thank you for being such an encouraging figure to me and for our long philosophical discussions that help me go through life with a little bit more wisdom.

Table of contents

Declaration of co-Authorship	iii
Abstract	v
Acknowledgments	vi
List of tables	xi
List of figures	xii
CHAPTER 1: General introduction	1
Introduction	2
Animal vocal signals: function and variation	2
Male-male interactions	4
Study site and study species.....	6
Thesis goals	8
Literature cited.....	9
Figures	13
CHAPTER 2: Diel and seasonal patterns in acoustic signalling of Savannah Sparrows	14
Chapter summary.....	15
Introduction	16
Material and methods.....	19
Study species and study site.....	19
Recording technique	20
Analysis of pairing and breeding status.....	20
Diel and seasonal variation	21
Statistical analysis.....	22
Results	23
Seasonal variation of song output.....	25
Song output variation with respect to breeding status.....	25
Dawn and dusk chorus, and daytime singing activity.....	24

Diel variation with respect to breeding status	23
Discussion.....	26
Social function of dawn chorus, afternoon singing activity, and dusk choruses..	26
Singing activity during breeding stages.	29
Literature cited	32
Figures	36
CHAPTER 3: Quiet violence: Songbirds respond to playback simulated rivals using soft songs as aggressive signals of intention	42
Chapter summary	43
Introduction.....	45
Material and methods	49
Study species and study site.....	49
Playback experiment	50
Playback stimuli.....	52
Analysis.....	53
Statistical analysis.....	55
Results	55
Initial period	56
One minute before attack	57
Discussion	57
Literature cited	62
Figures	68
CHAPTER 4: General discussion	71
Literature cited	76
Vita Auctoris	75

List of tables

Supplementary Table 2.1. Results of diel singing activity LMM standard least square comparisons for all interactions.	40
Supplementary Table 2.2. Results of seasonal singing activity LMM standard least square comparisons for all interactions.	41
Table 3.1. Summary of studies that experimentally investigated reliable predictors of attack in birds.	67

List of figures

Figure 1.1. Sound spectrogram showing the song of a Savannah Sparrow recorded at Bowdoin Scientific Station on Kent Island, New Brunswick, Canada, 2015.	13
Figure 2.1. Diel variation of the singing activity of Savannah Sparrows with respect to breeding stages.	36
Figure 2.2. Singing activity during the dawn chorus (30 min before sunrise) with respect to breeding stages (unpaired, fertile, incubation, hatchling and fledgling)	37
Figure 2.3. Singing activity during the daytime (time after sunrise and before sunset) with respect to five breeding stages (unpaired, fertile, incubation, hatchling and fledgling stages)	38
Figure 2.4. Singing activity during the dusk chorus (time after sunset) with respect to five breeding stages (unpaired, fertile, incubation, hatchling and fledgling stages).....	39
Figure 3.1. Diagram illustrating the playback experiment, based on the experimental design of Searcy et al. (2006).	68
Figure 3.2. Behaviors between attackers and non-attackers during the five-minute initial period of the playback experiment.	69
Figure 3.3. Behaviors between attackers and non-attackers during the one-minute prior to attack period of the playback experiment during taxidermy model presentation trials.	70

CHAPTER 1: General introduction

Introduction

Many animals produce vocal signals. These signals serve diverse functions, including mate attraction, territorial defense, alarm, identity recognition, and dominance signalling (Bradbury & Vehrencamp, 2011). Many animal signals vary seasonally, and this variation can inform our understanding of signal function and signalling behaviour (Slagsvold, 1977). Songbirds provide a compelling study system to investigate the function of acoustic signals because they have evolved complex and diverse vocal signals. In particular, songbirds use songs to defend territories and attract mates (Catchpole & Slater, 2008). My thesis focuses on investigating the functions of songs in Savannah Sparrows, *Passerculus sandwichensis*, with an emphasis on diel and seasonal variation of song output (Chapter 2), and the behaviour of birds during territorial interactions (Chapter 3). In this General Introduction, I provide a brief overview of the function of bird songs and seasonal variation in avian vocal behaviour and male vocal interactions, to provide background information for the two data chapters that follow.

Animal vocal signals: function and variation

Although animal vocal signals serve many functions including alarm, contact, and indication of food sources, two principal functions of vocal signals are territorial defense and mate attraction. For example, territorial male Galápagos Sea Lions, *Zalophus worllebaeki*, bark to attract females and to defend their harems from intruders (Kunc & Wolf, 2008). This dual function of animal vocal signals is well established, especially in

birds (i.e. the “dual function theory” of vocal signals by Catchpole & Slater 2008).

Indeed, male temperate songbirds are widely recognized to produce songs that are important both in territorial competition and mate attraction during the breeding season (reviewed in Catchpole & Slater, 2008).

Several lines of evidence reveal a strong association between vocal behaviour and mate attraction. Few experimental studies reveal that songs are important in mate choice but many observational studies show that unpaired males sing far more than paired males (e.g. Amrhein et al., 2004; Catchpole, 1973; Cuthill & Hindmarsh, 1985; Hennin et al., 2009). Once pairing or egg-laying has taken place, males also tend to reduce their song output (Amrhein et al., 2008; Ezaki, 1987). A common pattern observed during courtship behaviours is the increase in vocal activity or vocal complexity of male songs; in many temperate songbirds, song production peaks during the fertile period (Catchpole, 1973; Lampe & Espmark, 1987; Slagsvold, 1977). The investigation of the relationship between variation in singing activity and breeding stages provides us with a deeper understanding of the various functions of male songs.

In addition to the role that vocal signals play in mate attraction, a large body of evidence reveals that male vocal signals play a role in male-male interactions and territory defense. This was originally demonstrated through muting experiments, playback experiments, and speaker occupation experiments. Muting experiments demonstrated that muted males were unable to maintain their territories (e.g. Red-winged Blackbirds, *Agelaius phoeniceus*: Peek, 1972; Scott’s Seaside Sparrows,

Ammodramus maritimus peninsulae: McDonald, 1989). Playback experiments established that the playback of male songs triggered aggression from nearby neighbours (e.g. Indigo Bunting, *Passerina cyanea*: Emlen, 1971, Song Sparrows, *Melospiza melodia*: Searcy et al. 2006); and speaker occupation experiments confirmed that playback of songs effectively repelled rival males from speaker-occupied territories (Great Tits, *Parus major*: Krebs et al., 1978; Song Sparrows: Nowicki et al., 1998; White-throated Sparrows: Falls, 1988). Songs are thus important signals for maintaining territories, and effective signals for deterring rivals. Exploring how males use their songs during male-male interactions provides insight into the role of songs in territory establishment and maintenance of boundaries.

Male-male interactions

Territorial interactions between rival males routinely involve aggression (Todt & Naguib, 2000). Given that aggressive encounters may involve costly physical attacks, many animals have evolved agonistic signals to facilitate communication without physical confrontation during aggressive contests (Johnstone, 1997). As contests escalate, animal signals can transition from low-level threat signals to high-level threat signals. For example, Sac-winged Bats, *Saccopteryx bilineata*, call at lower pitches and greater rates when territorial contests escalate (Behr et al. 2009) and Green Frogs, *Rana clamitans*, decrease the pitch of their calls when territorial contests escalate (Bee et al., 2000; Reichert & Gerhardt, 2013). In addition, songbirds often change their singing behaviour to escalate or de-escalate contests (Todt & Naguib, 2000a). Previous research

on signal escalation has focused on song overlapping, song-type matching, song switching, or modulation of the fine structure of song elements (e.g. Burt et al., 2001; DuBois et al., 2009; Naguib, 2005; Searcy et al., 2006), and many of these signalling behaviours change as contests escalate or de-escalate.

The best-studied system of territorial contest escalation comes from Song Sparrows (Akçay et al., 2013; Beecher & Campbell, 2005; Searcy & Beecher, 2009). In a series of experiments designed to study aggressive signalling during Song Sparrow territorial disputes, Searcy and Beecher (2009) established a compelling model for contest escalation. Their hierarchical signalling model proposes that during a dyadic interaction, a male may escalate a contest by singing the same song as his rival's songs, a behaviour known as song matching, or de-escalate by switching to a non-matching song (Akçay et al. 2013). Similar investigations have been conducted in other taxa, which all point towards the possible existence of graded signals of contest escalation (e.g. Hof & Podos 2013). Songbirds can use other vocal communication strategies to escalate interactions such as song frequency matching (Baker et al., 2012), increases in song frequency and song rate (DuBois et al., 2009), decreases in song amplitude (Ballentine et al., 2008; Grieves et al., 2015; Hof & Hazlett, 2010; Reichard et al., 2011; Ritschard et al., 2012; Searcy et al., 2006; Xia et al., 2013), and production of non-song calls (Ballentine et al., 2008, Baker et al., 2012, Elie & Theunissen 2015). Remarkably few studies have explored the precise function of signals during aggressive interactions or described the graded signalling strategies that comprise a hierarchical signalling model. The second data chapter of my thesis explores these ideas.

Researchers today focus on expanding our understanding of vocal signals to include both songs (i.e. complex learned vocalizations, usually produced by males during the breeding season) as well as calls (i.e. simple, short non-learned vocalizations, sometimes produced by both males and females throughout the year; e.g. Gill et al., 2015; Elie & Theunissen, 2015). Past research focused on songs as the main signal that plays a territorial and attractive function, whereas the importance of calls during these interactions remained poorly studied. Like songs, calls are often produced during periods of territorial defence and mate attraction, and have been shown to be produced in diverse contexts (Marler, 2004) and in large numbers (Beckers & Gahr, 2010). The second chapter of my thesis explore calls as potential signalling strategies to escalate interactions.

Study site and study species

My thesis research took place at the Bowdoin Scientific Station on Kent Island, New Brunswick, Canada (44°35'N 66°46'W), a 200-acre island in the Bay of Fundy. This island is home to a population of Savannah Sparrows with strong philopatry (Wheelwright & Mauck, 1998). Savannah Sparrows migrate from their wintering grounds in the southern United States (Woodworth et al. 2015) to their breeding sites across the United States and Canada (Bédard & LaPointe, 1984; Tufts, 1986). The birds on Kent Island have been studied annually for many decades (Williams et al. 2013). Each year, more than one hundred Savannah Sparrows at the study site are marked with a unique combination of leg bands to facilitate recognition of individuals in the field. Male

Savannah Sparrows arrive from migration in mid-April and begin to defend breeding territories from rival males (Woodworth et al. 2015). This is an important moment in the annual cycle of males; if they are to attract females and breed successfully, they need to establish a breeding territory (Potter, 1972). During territorial establishment, male Savannah Sparrows interact with other males by singing a single song type (Figure 1.1). Males appear to learn their songs within the first eight months of life (Wheelwright et al., 2008) and songs remain constant throughout an individual's life (Williams et al., 2013).

During territory establishment, male Savannah Sparrows use an array of vocal and visual signals during aggressive interactions with rivals. Male song is a regular component of male behaviour during territory establishment (Wheelwright & Rising, 2008). In addition to their songs, males emit a variety of calls during territorial encounters, including aggressive flight calls and hostile notes (a.k.a. buzz calls) (Gobeil, 1970; Wheelwright & Rising, 2008). Males also engage in agonistic visual displays that include parallel walking, flutter flights, tail raising, bill gaping, ptiloerection of crest feathers, and bouts of physical chases (Potter, 1972; Wheelwright & Rising, 2008). The relative importance of these diverse vocalizations and physical displays are poorly studied; they may be critical for territorial signalling in Savannah Sparrows, and they may comprise a graded system of aggressive signalling, although this has never been studied previously.

Thesis goals

In this thesis, I explore the vocal behaviour of Savannah Sparrows with a focus on communication in the context of territorial aggression and mate attraction. My thesis has two components. Chapter 2 is an observational study based on detailed recordings of individually-marked Savannah Sparrows. I describe diel and seasonal variation in vocal output, evaluate how male singing behaviour changes with different breeding stages, and conduct a detailed analysis of recordings of territorial males. In Chapter 2, I ask the question: *what is the pattern of diel and seasonal variation in vocal activity of Savannah Sparrows, and what does this reveal about the functions of their vocal signals?* Chapter 3 is an experimental study based on a recently developed playback protocol that has been used to study aggressive signalling in other songbirds. I conducted a playback and model-presentation experiment to study the graded signals that are associated with contest escalation during Savannah Sparrow territorial encounters. In Chapter 3, I ask the question: *do Savannah Sparrows produce signals that predict physical attack during territorial interactions?* With this body of research, I hope to expand our understanding of vocal behaviour of Savannah Sparrows specifically, and, more broadly, to develop a better understanding of seasonal and diel variation in vocal signals and the dynamics of signalling behaviour during aggressive territorial encounters.

Literature cited

- Akçay, Ç., Tom, M. E., Campbell, S. E., & Beecher, M. D. (2013). Song type matching is an honest early threat signal in a hierarchical animal communication system. *Proceedings of the Royal Society B: Biological Sciences*, 280, 20122517.
- Amrhein, V., Johannessen, L. E., Kristiansen, L., & Slagsvold, T. (2008). Reproductive strategy and singing activity: blue tit and great tit compared. *Behavioral Ecology and Sociobiology*, 62, 1633–1641.
- Amrhein, V., Kunc, H. P., & Naguib, M. (2004). Seasonal patterns of singing activity vary with time of day in the nightingale (*Luscinia megarhynchos*). *The Auk*, 121, 110–117.
- Baker, T. M., Wilson, D. R., & Mennill, D. J. (2012). Vocal signals predict attack during aggressive interactions in black-capped chickadees. *Animal Behaviour*, 84, 965–974.
- Ballentine, B., Searcy, W., & Nowicki, S. (2008). Reliable aggressive signaling in swamp sparrows. *Animal Behaviour*, 75, 693–703.
- Beckers, G. J. L., & Gahr, M. (2010). Neural Processing of Short-Term Recurrence in Songbird Vocal Communication. *PLOS ONE*, 5, e11129.
- Bédard, J., & LaPointe, G. (1984). Banding returns, arrival times, and site fidelity in the Savannah Sparrow. *The Wilson Bulletin*, 96, 196–205.
- Bee, M. a, Perrill, S. a, & Owen, P. C. (2000). Male green frogs lower the pitch of acoustic signals in defense of territories: a possible dishonest signal of size? *Behavioral Ecology*, 11, 169–177.
- Beecher, M. D., & Campbell, S. E. (2005). The role of unshared songs in singing interactions between neighbouring song sparrows. *Animal Behaviour*, 70, 1297–1304.
- Behr, O., Knörnschild, M., & Von Helversen, O. (2009). Territorial counter-singing in male sac-winged bats *Saccopteryx bilineata*: Low-frequency songs trigger a stronger response. *Behavioral Ecology and Sociobiology*, 63, 433–442.
- Bradbury, J. W., & Vehrencamp, S. L. (1998). *Principles of animal communication*. Sunderland. Sinauer Associates.
- Burt, J. M., Campbell, S. E., & Beecher, M. D. (2001). Song type matching as threat: a test using interactive playback. *Animal Behaviour*, 62, s163–1170.

- Catchpole, C. K. (1973). The Functions of advertising song in the Sedge Warbler (*Acrocephalus schoenobaenus*) and the Reed Warbler (*A. scirpaceus*). *Behaviour*, 46, 300–319.
- Catchpole, C. K., & Slater, P. J. B. (2008). *Bird song: Biological themes and variations*.
- Cuthill, I., & Hindmarsh, A. (1985). Increase in starling song activity with removal of mate. *Animal Behaviour*, 33, 326–328.
- DuBois, A. L., Nowicki, S., & Searcy, W. A. (2009). Swamp sparrows modulate vocal performance in an aggressive context. *Biology Letters*, 5, 163–165.
- Emlen, S. T. (1971). The role of song in individual recognition in the indigo bunting. *Ethology*, 28, 241–246.
- Ezaki, Y. (1987). Male time budgets and recovery of singing rate after pairing in polygamous great reed warblers. *Japanese Journal of Ornithology*, 36, 1–11.
- Fallow, P. M., Gardner, J. L., & Magrath, R. D. (2011). Sound familiar? Acoustic similarity provokes responses to unfamiliar heterospecific alarm calls. *Behavioral Ecology*, 22, 401–410.
- Falls, J. B. (1988). Does song deter territorial intrusion in White-throated Sparrows (*Zonotrichia albicollis*)? *Canadian Journal of Zoology*, 66, 206–211.
- Gobeil, R. E. (1970). Vocalizations of the Savannah Sparrow. *Bird-Banding*, 41, 18–21.
- Grieves, L. A., Logue, D. M., & Quinn, J. S. (2015). Ready to fight: Reliable predictors of attack in a cooperatively breeding, non-passerine bird. *Ethology*, 121, 1154–1165.
- Hennin, H. L., Barker, N. K. S., Bradley, D. W., & Mennill, D. J. (2009). Bachelor and paired male rufous-and-white wrens use different singing strategies. *Behavioral Ecology and Sociobiology*, 64, 151–159.
- Hof, D., & Hazlett, N. (2010). Low-amplitude song predicts attack in a North American wood warbler. *Animal Behaviour*, 80, 821–828.
- Johnstone, R. A. (1997). The evolution of animal signals. *Behavioural Ecology: An Evolutionary Approach*, 155–178.
- Krebs, J., Ashcroft, R., & Webber, M. (1978). Song repertoires and territory defence in the great tit. *Nature*, 271, 539–542.
- Kunc, H. P., & Wolf, J. B. W. (2008). Seasonal changes of vocal rates and their relation to territorial status in male Galápagos sea lions (*Zalophus worrebaeki*). *Ethology*, 114, 381–388.

- Lampe, H. M., & Espmark, Y. O. (1987). Singing activity and song pattern of the Redwing *Turdus iliacus* during the breeding season. *Ornis Scandinavica*, 18, 179–185.
- Marler, P. (2004). Bird calls: their potential for behavioral neurobiology. *Annals of the New York Academy of Sciences*, 1016, 31–44.
- McDonald, M. V. (1989). Function of song in Scott's seaside sparrow, *Ammodramus maritimus peninsulae*. *Animal Behaviour*, 38, 468–485.
- Naguib, M. (2005). Singing interactions in songbirds: implications for social relations and territorial settlement. *Animal Communication Networks*, 300–319.
- Nowicki, S., Searcy, W. A., & Hughes, M. (1998). The territory defense function of song in song sparrows: a test with the speaker occupation design. *Behaviour*, 135, 615–628.
- Peek, F. W. (1972). An experimental study of the territorial function of vocal and visual display in the male red-winged blackbird (*Agelaius phoeniceus*). *Animal Behaviour*, 20, 112–118.
- Potter, P. E. (1972). Territorial behavior in Savannah Sparrows in southeastern Michigan. *The Wilson Bulletin*, 84, 48–59.
- Reichard, D. G., Rice, R. J., Vanderbilt, C. C., & Ketterson, E. D. (2011). Deciphering information encoded in birdsong: male songbirds with fertile mates respond most strongly to complex, low-amplitude songs used in courtship. *The American Naturalist*, 178, 478–487.
- Reichert, M. S., & Gerhardt, H. C. (2013). Gray tree frogs, *Hyla versicolor*, give lower-frequency aggressive calls in more escalated contests. *Behavioral Ecology and Sociobiology*, 67, 795–804.
- Ritschard, M., van Oers, K., Naguib, M., & Brumm, H. (2012). Song amplitude of rival males modulates the territorial behaviour of great tits during the fertile period of their mates. *Ethology*, 118, 197–202.
- Searcy, W. a., Anderson, R. C., & Nowicki, S. (2006). Bird song as a signal of aggressive intent. *Behavioral Ecology and Sociobiology*, 60, 234–241.
- Searcy, W., & Beecher, M. D. (2009). Song as an aggressive signal in songbirds. *Animal Behaviour*. *Animal Behaviour*, 78, 1281-1292.
- Slagsvold, T. (1977). Bird song activity in relation to breeding cycle, spring weather, and environmental phenology. *Ornis Scandinavica*, 8, 197–222.
- Todt, D., & Naguib, M. (2000). Vocal interactions in birds: The use of song as a model in communication. *Advances in the Study of Behavior*, 29, 247-296.

- Tufts, R. W. (1986). Birds of Nova Scotia. (3rd ed). Halifax, NS: Nimbus Publ. Ltd. and Nova Scotia Mus.
- Wheelwright, N., & Rising, J. (2008). Savannah Sparrow (*Passerculus sandwichensis*). Number 45. The Birds of North America. Cornell University, Laboratory of Ornithology, Ithaca, New York, USA.
- Wheelwright, N. T., & Mauck, R. A. (1998). Philopatry, natal dispersal, and inbreeding avoidance in an island population of Savannah Sparrows. *Ecology*, 79, 755–767.
- Wheelwright, N. T., Swett, M. B., Levin, I. I., Kroodsma, D. E., Freeman-Gallant, C. R., & Williams, H. (2008). The influence of different tutor types on song learning in a natural bird population. *Animal Behaviour*, 75, 1479–1493.
- Williams, H., Levin, I. I., Norris, D. R., Newman, A. E. M., & Wheelwright, N. T. (2013). Three decades of cultural evolution in Savannah sparrow songs. *Animal Behaviour*, 85, 213–223.
- Woodworth, B. K., Mitchell, G. W., Norris, D. R., Francis, C. M., & Taylor, P. D. (2015). Patterns and correlates of songbird movements at an ecological barrier during autumn migration assessed using landscape-and regional-scale automated radiotelemetry. *Ibis*, 157, 326–339.
- Xia, C., Liu, J., Alström, P., Wu, Q., & Zhang, Y. (2013). Is the soft song of the brownish-flanked bush warbler an aggressive signal? *Ethology*, 119, 653–661.

Figures




Figure 1.1. Sound spectrogram showing the song of a Savannah Sparrow recorded at Bowdoin Scientific Station on Kent Island, New Brunswick, Canada, 2015.

CHAPTER 2: Diel and seasonal patterns of acoustic signalling in Savannah

Sparrows

Chapter summary

Animal vocal output can change daily and seasonally. Diel and seasonal variation in vocal output can be an indicator of signal function and studies of daily and seasonal rhythms can provide us with deeper insight into animal behaviour. In this study, we quantified diel and seasonal variation in song output in a breeding population of Savannah Sparrows, *Passerculus sandwichensis*. We used automated recorders to collect extensive recordings across the breeding season. We found significant diel and seasonal variation in vocal output with respect to male pairing and breeding status. In terms of diel variation, males showed a peak in the early morning, consistent with dawn chorus behaviour, but this pattern was common only prior to pairing. After pairing, males showed a peak in the evening, consistent with dusk chorus behaviour. In terms of seasonal variation, Savannah Sparrows showed the highest level of song output in May and the lowest level in August. Males showed the highest singing activity early in the year, when birds were first establishing territories and also when females were incubating. Unpaired territorial males had the most pronounced dawn chorus and higher levels of morning singing activity. A dusk chorus occurred later in the breeding season, during the fertile, incubation, hatchling, and fledgling periods, with paired males participating most heavily. These patterns suggest that dawn choruses have a territorial function whereas dusk choruses may have a female-related function in this species. We conclude that Savannah Sparrows show dynamic patterns of diel and seasonal variation, where vocal output varies with social context at different stages of their breeding season.

Introduction

Many territorial animals use vocal signals that vary daily and seasonally.

Variation in signalling output is often linked with changes in seasons, when the levels of abiotic factors such as light and temperatures change. Several hypotheses have been proposed to explain why animals vary their vocalizations daily and seasonally and these fall under three main categories: temporal variation in vocal output can be related to mechanistic processes (e.g. the hormones that influence song output), social processes (e.g. the social contexts that affect vocal output), or environmental features (e.g. the environmental constraints on vocal output) (Burt & Vehrencamp, 2005; Cuthill & Macdonald, 1990; Kacelnik & Krebs, 1983; Mace, 1987; Staicer et al., 1996). The influence of social forces such as pairing status (Catchpole, 1973; Cuthill & Hindmarsh, 1985; Demko et al., 2013) and breeding status (Bruni & Foote, 2014; Foote & Barber, 2009; Hanski & Laurila, 1993; Zhang, Celis-Murillo, & Ward, 2016) have a profound influence on vocal output, and understanding the connection between social activities and changes in vocal behaviour has proven to be a revealing area of research. In this study, we investigate the social processes that influence the diel and seasonal vocal variation in a North American songbird species, Savannah Sparrows, *Passerculus sandwichensis*.

The singing activity of many animals varies with time of day, especially in species that exhibit dawn and dusk choruses (reviewed in Staicer et al. 1996). Dawn and dusk choruses are periods of heightened vocal activity that appear to be important for

territorial defense, mate attraction, and extra-pair mate attraction (Burt & Vehrencamp, 2005; Catchpole, 1973; Kunc, Amrhein, & Naguib, 2005; Poesel et al., 2006; Slagsvold, Dale, & Sætre, 1994). Dawn choruses usually start before sunrise and conclude at sunrise, or shortly after sunrise, depending on the species or time of the year (reviewed in Staicer et al. 1996). Similarly, dusk choruses begin at sunset, and continue after sunset (e.g. Cuthill & Macdonald, 1990; Lein, 2007). Animals that show dawn and dusk choruses include primates (Schel & Zuberbühler 2011), lizards (Ord, 2008), and birds (Burt & Vehrencamp, 2005). Songbirds are particularly well-known for their dawn and dusk choruses, and they provide a model system in the field of acoustic communication to study diel variation in vocal output.

In addition to diel variation in vocal output, the singing activity of animals can also vary with pairing and breeding status. In many species of songbirds, for example, males sing extensively until they are paired and singing activity is greatly reduced after pairing (e.g. Sedge Warblers, *Acrocephalus schoenobaenus*: Catchpole, 1973), although other species continue singing even after they are paired (e.g. Reed Warblers, *Acrocephalus scirpaceus*: Catchpole, 1973). In some species, males use different types of signals when unpaired versus paired (Catchpole, 1973) or modulate their songs with different breeding stages (Zhang et al., 2016). The singing activity of songbirds can also vary with female breeding stages, often showing levels of high output during the fertile period (Amrhein et al., 2008; Foote & Barber, 2009; Slagsvold, 1977) or during both the fertile and incubation periods (e.g. Bruni & Foote, 2014). In other species, males increase their song rate during the nest-building stage (e.g. Northern Mockingbirds,

Mimus polyglottos: Logan, 1983) or vary their song complexity during periods of female attraction (e.g. Blue Grosbeaks, *Passerina caerulea*: Ballentine et al., 2003). Singing activity can also vary during the egg-laying period. In Nightingales, *Luscinia megarhynchos*, for example, males lower their song output and reduce their song overlapping during the egg-laying period (Amrhein et al. 2004). Investigating variation in song output provides an opportunity for understanding the role of male songs with their association to female breeding status.

In this study, we collected long-term recordings to quantify seasonal and diel output in Savannah Sparrows, and we compare vocal output to breeding activities and observational data. We tested the hypotheses that vocal output in song varies with time of day and with time of year. We made *a priori* predictions about the relationship between variation in vocal output and breeding activities. We predicted that if the main function of dawn and morning chorus singing in male Savannah Sparrows is to defend and maintain their territory, then the song output should not vary with female breeding stages. Alternatively, if the main function of the dawn chorus is related to female breeding status, then song output should vary throughout the breeding cycle and serve different functions.

Material and methods

Study species and study site

Savannah Sparrows are small migratory birds that live in open grasslands throughout North America (Bédard & LaPointe, 1984). Males migrate from their wintering grounds in the southern United States to their breeding sites across the United States and Canada (Bédard & LaPointe, 1984; Tufts, 1986). We conducted our research at the Bowdoin Scientific Station on Kent Island, New Brunswick, Canada (44°35'N 66°46'W), a 200-acre island in the Bay of Fundy. This island is home to a population of Savannah Sparrows with strong philopatry (Wheelwright & Mauck, 1998). The birds in this population have been studied for more than three decades (Williams et al., 2013). Each year, more than one hundred Savannah Sparrows at the study site are marked with a unique combination of leg bands to facilitate recognition of individuals in the field. Male Savannah Sparrows breeding on Kent Island arrive from migration in mid-April, about one month before females' arrival in mid-May, and begin to defend breeding territories from rival males (Woodworth et al., 2015). During territorial establishment, male Savannah Sparrows interact with other males by singing a single song type. This is an important period for males; if they can establish a breeding territory they will attract females more easily (Potter, 1972). Kent Island males' territories size average is 0.21 ha and range in size between 0.03-0.60 ha (Wheelwright & Rising, 2008) which allow easy monitoring of multiple interactions between multiple males.

Recording technique

In 2014, we recorded 34 males using automated recorders, Wildlife Acoustics Song Meters (model: SM2; see Mennill et al. 2012). Given the small territories of Savannah Sparrows, these devices allowed us to record between one and seven males simultaneously with a single recorder. We recorded males for ten 24h-long recording sessions between mid-April and mid-September, with 14-day intervals between subsequent recording sessions at each site. In all recordings where multiple males were present, we distinguished individuals on the basis of their individually distinctive song (see Wheelwright et al., 2008).

Analysis of pairing and breeding status

To determine breeding stages of the 34 males in our diel and seasonal variation analysis, we collected behavioural observations and monitored nests every two days. Following Foote & Barber (2009), we divided the breeding season into five different periods from May to September: (1) males that were unpaired (the period starting from males' arrival around mid-April, up to 15 days later, start of May; females were absent during this period), (2) males paired with a fertile female (the period corresponding to 8 days prior to the laying of the penultimate egg), (3) males with a female incubating eggs (a period of 12 days), (4) males with a female provisioning nestlings (a period of 9 days), and (5) males with a female provisioning fledglings (a period starting from the end of the nestling stage to the start of the next brood or if no consecutive brood was attempted, a period of 7 days after the end of nestling period). We estimated the length

of the fertile period based on the following logic: female Savannah Sparrows are thought to be fertile during the 1-3 days when nests are built and during the following period of 3-5 days when eggs are laid (1egg/day). We calculated the start of each breeding period based on the fertile period. But, in some cases, we estimated the first day of incubation by back-dating 12 days from known hatch dates because in those cases the nests were found later during the incubation stage thus the first day of egg-laying could not be determined (Dixon, 1978; Wheelwright & Rising, 2008).

Diel and seasonal variation

In order to compare song output between dawn chorus, daytime, and dusk chorus, we defined the dawn chorus as the period 30 minutes before sunrise, a time period that included some of the civil twilight and dawn (as in Liu, 2004; Naguib et al., 2016). Daytime was defined as the period starting from sunrise to sunset. Dusk chorus was defined as the period starting from sunset up to the last songs. Finally, for each of the five breeding periods, we investigated diel variation by calculating and comparing hourly values for the total song output during dawn chorus, daytime, and dusk chorus. Sunrise and sunset times for each day were obtained from the National Research Council, Herzberg Institute of Astrophysics sunrise/sunset calculator (www.nrc-cnrc.gc.ca). We excluded days where weather affected recordings (i.e. very rainy or windy days). On average, we included 7 ± 3 days of recording across the season for each individual.

Statistical analysis

We conducted an analysis of diel variation in song output using linear mixed models to compare diel song output between breeding stages. We included time of day (subdivided into 1-hr periods), breeding stage (unpaired, fertile, incubation, hatchling, or fledgling), and interaction between time and breeding stage as our fixed factors. We included individuals as a random effect to account for the fact that the same males were sampled repeatedly.

Finally, we compared the difference in song output between breeding stages for three periods: the dawn chorus, daytime, and the dusk chorus. Because the distribution of our data was non-normal, we compared song output across those time periods using a Wilcoxon test. To determine how song output varies between breeding stages, we then compared the average number of songs between breeding stages using a Wilcoxon test.

We analyzed the seasonal variation of the singing activity of Savannah Sparrows using linear mixed models to determine the period with highest and lowest song output per hour. For the analysis of seasonal variation, we included Julian day (83 days between 4 May and 26 July 2014), breeding stage (unpaired, fertile, incubation, hatchling, or fledgling) and the interaction between Julian day and breeding stage as our fixed factors. We included male identity as a random effect to account for the fact that the same males were sampled repeatedly.

We used JMP v12 (SAS Institute Inc. 2015) for all our statistical analyses. We considered results with $P < 0.05$ significant and we represented values as means \pm SE. Time of sunrise and sunset from the start of our recording on 4 May until the end of our recordings on 26 July 2014 varied respectively from 07:05 h to 06:06 h for sunrise and from 19:55 h to 21:00 h for sunset.

Results

Diel variation with respect to breeding status

We conducted an analysis of diel variation in song output using linear mixed models to compare diel song output between breeding stages. Savannah Sparrow song output varied significantly with respect to breeding stages ($F_{4,4687} = 13.9$, $P < 0.001$), with respect to time of day ($F_{1,4698} = 22.5$, $P < 0.001$), and interaction between breeding stage and time of day ($F_{4,4698} = 52.4$, $P < 0.001$), ($n = 34$ males) (Fig. 2.1; Supplementary Table 2.1). Savannah Sparrows' diel song output showed significant variation between time of day in the unpaired period ($t_{83} = 8.9$, $P < 0.001$), the fledgling period ($t_{4641} = 6.2$, $P < 0.001$), and incubation period ($t_{4670} = 5.9$, $P < 0.001$) ($n = 30$ males). The diel song output during the fertile period ($t_{4724} = 0.42$, $P = 0.67$) and the hatchling period ($t_{4720} = 0.84$, $P = 0.39$) did not vary significantly across time of day (Fig. 2.1). We obtained a R^2 value of 0.12 and the distribution of values of random effect had a variance of 0.07.

One striking difference between unpaired males and males at other breeding stages (fertile, incubation, hatchling, fledgling) was the difference in song output in the

dawn, daytime, and dusk chorus song activities. The song output of unpaired males peaked in the morning around 07:00 h and then males reduced their singing activity from late morning into the evening (between approximately 11:00 h to 22:00 h). Males at other breeding stages had a small increase in singing activity early morning (around 5:00 h), maintained their singing activity in morning (between 06:00 h to 12:00 h) and the afternoon evening (between approximately 12:00 h to 20:00 h) and increased their singing activity during dusk chorus (between approximately 20:00 h to 22:00 h).

Dawn chorus, dusk chorus, and daytime singing activity

We compared the difference in song output between breeding stages for three periods: the dawn chorus, daytime, and the dusk chorus. When comparing different breeding stages across three daytime periods (morning chorus, daytime, and dusk chorus), the total song output during the dawn chorus (30 min before sunrise until sunrise) was significantly higher when males were unpaired than all other breeding stages (Fig. 2.3), including the fertile period ($Z=3.02$, $P=0.003$), incubation period ($Z=2.61$, $P=0.009$), hatchling period ($Z=2.33$, $P=0.02$), and fledgling period ($Z=2.20$, $p=0.03$). Once males were paired, their singing activity during the dawn chorus decreased by about half compared to the unpaired stage and did not differ significantly between stages. During the day, male singing activity did not significantly differ between breeding stages except between the unpaired and fledgling stage ($Z=2.26$, $P=0.02$), where unpaired males sang more than males in the fledgling stage (Figure 2.4). During the dusk chorus period, unpaired males sang significantly less than any other breeding

stage (Fig. 2.5). Unpaired males sang significantly less compared to the fertile period ($Z=-4.24$, $P<0.0001$), the incubation period ($Z=-2.79$, $P=0.005$), the hatchling period ($Z=-2.04$, $P=0.04$), and the fledgling period ($Z=-2.57$, $P=0.01$). The song output during the dusk chorus was also significantly smaller during the hatchling stage compared to the fertile stage ($Z=-2.26$, $P=0.02$).

Song output variation with respect to breeding status

Song output during the incubation stage was significantly higher than any other stage except the unpaired stage ($Z=-6.7$, $P<0.001$) and was not significantly different from the hatchling stage ($Z=0.54$, $P=0.58$). Song output per hour during the fledgling stage was significantly lower than the unpaired stage ($Z=-2.67$, $P=0.008$), and the hatchling stage ($Z=2.02$, $P=0.04$) but not the fertile stage ($Z=-0.93$, $P=0.35$).

Seasonal variation of song output

We analyzed the seasonal variation of the singing activity of Savannah Sparrows using linear mixed models to determine the period of the breeding year with highest and lowest song output per hour. The song output of Savannah Sparrows varied seasonally (Linear mixed model $F_{1, 4372}=10.91$, $P=0.001$), with respect to breeding stages ($F_{4, 4627}=6.4$, $P<0.0001$), and interaction between breeding stage and julian day ($F_{4, 4680}=3.9$, $P=0.003$) (Supplementary Table 2.2). Males decreased their song output per hour throughout the breeding season. Savannah Sparrows sang the most per hour early in

the breeding season (May and June). We obtained a R^2 value of 0.08 and the distribution of values of random effect had a variance of 0.077

Discussion

Our study of diel and seasonal variation in song output of Savannah Sparrows highlights clear patterns of singing activity throughout the day and the breeding season and sheds light on the function of songs in species. Males showed diel variation in their song output with respect to pairing status of males and breeding status of females. Unpaired males sang most during the dawn chorus compared to paired males in all breeding stages, whereas unpaired males sang least during the dusk chorus compared to paired males in all other breeding stages. In addition, males showed vocal variation with respect to breeding stages. Males with fledglings sang the least compared to males during all other breeding stages, while singing activity was highest during the incubation stage and unpaired stage. Male singing activity also varied seasonally. Males sang with the highest song output per hour early in May, when they were unpaired and establishing territories. Later in the breeding season in June, the average male singing activity increased again although less than in the unpaired stage.

Social function of dawn chorus, daytime singing activity, and dusk choruses

Variation in vocal activity can reflect important aspects of an animal's life history. In our study, the variation in the singing activity of Savannah Sparrows supports the idea that songs have a territorial function early in the breeding season and early in the

morning. The high singing activity in the mornings of unpaired Savannah Sparrows, when females are absent from the island altogether, strongly suggests that the morning is a period of importance for territorial interactions to settle and defend territories in the month of April and May. Indeed, if the dawn chorus had an attractive function, unpaired males would not be expected to sing during this period. Defending territories with songs in Savannah Sparrows is a crucial activity that increases males' pairing success (Reid & Weatherhead, 1990). Particularly, pairing success in Savannah Sparrows appears to be influenced by male territory size or male song rate in this species (Reid & Weatherhead 1990). This may explain why males arrive at their breeding site about one month before females and sing and fight intensively for a month. This remarkable time difference between female and male migration arrival reveals the importance of establishing and maintaining a territory in this species.

Once paired, Savannah Sparrows' morning singing activity decreased by about half and remained constant throughout the breeding stages. This suggests that dawn chorus still plays an important social role even after territorial establishment. Morning choruses after pairing in temperate birds have been suggested to act as a social period during which males maintain their territories (Amrhein & Erne, 2006; Erne & Amrhein, 2008; Foote et al., 2011), strengthen their pair bond (Erne & Amrhein, 2008) or solicit extra-pair copulations (Poesel et al. 2006). For example, the dawn chorus is an honest signal of male quality in Eastern Kingbirds, *Tyrannus tyrannus* (Murphy et al., 2008). When singing more and earlier, male Eastern Kingbirds have a higher chance of having extra-pair copulations. Similar to Savannah Sparrows, other bird species, like the

Collared Flycatcher, *Ficedula albicollis*, the Chiffchaff, *Phylloscopus collybita*, and the Nightingale, *Luscinia megarhynchos* (Kunc et al. 2005) maintain a dawn chorus throughout the breeding stages and show that dawn chorus is used for territory defence and maintenance only (Kunc et al., 2005; Pärt, 1991; Rodrigues, 1996; Staicer et al., 1996).

Our study provides quantitative evidence that Savannah Sparrows produce a dusk chorus. Dusk choruses showed striking patterns of seasonal variation, where males sang little or no dusk chorus in the absence of females, but once paired, males sang pronounced dusk choruses. This pattern suggests that in this species, the dusk chorus may function as a period of time during which males target females although males may also use this period for territorial purposes too. Dusk chorus has been hypothesized to function as a period during which males defend territories, obtain extra pair copulations or maintain their female-male pair bond in other species (Cuthill & Macdonald, 1990; Erne & Amrhein, 2008). Future observational studies and playback experiments on dusk chorus songs could shed light on the function of songs during the dusk chorus songs in Savannah Sparrows. Comparing the song output between unpaired males during the breeding season (i.e. from May till August) with paired males could answer whether dusk songs are sung only by paired males, in which case dusk chorus may function as a period of time during which paired males target females only.

Singing activity during breeding stages

In our study, we showed that singing activity changes as the breeding cycle progresses. When excluding the unpaired stage, Savannah Sparrows sang the highest when males were paired with an incubating female. Researchers have argued that an increase in song output during the incubation stage acts as a mate guarding, a female stimulant to incubate (Foote & Barber, 2009), or an all clear signal to indicate to females that the male is nearby and that no predators are around (Johnson & Kermott 1991). In the case of Savannah Sparrows, incubation may be an important period for males to obtain extra paired copulations. Similar to our study, Slagsvold's research (1977) on seasonal variation in the song behaviour of 20 species of songbirds showed that song activities are tightly related to breeding cycles and that singing activity peaks within few days of egg-laying. This finding was supported by other studies in the European Redwings, *Turdus iliacus* (Lampe & Espmark, 1987) and *Acrocephalus* warblers (Catchpole, 1973). Although, it is generally supported that singing activity peaks within the fertile period of the females (a period including egg-laying and nest-building) as suggested by the fertility announcement hypothesis (Møller, 1991), there are numerous cases of songbirds similar to Savannah Sparrows that do not follow this pattern and do not support this hypothesis. For example, Willow Warblers, *Phylloscopus trochilus*, do not sing during fertile periods (Gil, Graves, & Slater, 1999), and the songs of Song Sparrows, *Melospiza melodia*, peak during incubation (Foote & Barber, 2009). In addition, singing during the incubation near or on the nest is not uncommon and was

shown to occur in about 160 species in North America (Leonard, 2008) and birds like the Common Chaffinch, *Fringilla coelebs* (Hanski & Laurila, 1993, Rodrigues 1996), and Willow Warbler, *Phylloscopus trochilus* (Gil et al. 1999).

Singing activity during the

Finally, in our study male Savannah Sparrows showed a high degree of variation in terms of singing activity throughout the breeding season. According to Amrhein et al (2008), this seasonal variation pattern in song output can partly be influenced by their reproductive strategies. Amrhein et al. (2008) compared both the singing activity pattern and reproductive strategy of two different tits, Blue Tits, *Cyanistes caeruleus*, and Great Tits, *Parus major*. Blue Tits, which are a facultatively polygamous and have a short breeding period and do not produce second broods; whereas Great Tits are a socially monogamous species and can have second broods (Amrhein et al. 2008). Great tits were found to increase their song output during egg laying and incubation periods and continued singing later in the breeding season, while Blue Tits decreased their song output after incubation of their first brood. Amrhein et al. (2008) explain that the multi-brooded reproductive strategy may constrain Great Tits to defend their territories over longer periods of time. Similar to Great Tits, Savannah Sparrows may be constrained to sing over longer periods of time (from April till July) due to their multi-brooded reproductive strategy that encourages males to sing (Wheelwright, & Rising, 2008).

In conclusion, we showed that singing activity varies throughout the day and breeding season, and with respect to breeding stages in this species. In terms of diel

variation, unpaired territorial males had the most pronounced dawn chorus and higher levels of morning singing activity. A dusk chorus occurred later in the breeding season, during the fertile, incubation, hatchling, and fledgling periods. In addition, Savannah Sparrows singing activity is highest during the month of May and June. These patterns suggest that Savannah Sparrows show dynamic patterns of diel and seasonal variation, where vocal output varies with social context at different stages of their breeding season. Our study highlights the general singing patterns of Savannah Sparrows throughout the breeding season. Analysis of seasonal variation of singing like our study can provide insight into communication systems of song birds that heavily rely on singing interactions.

Literature cited

- Amrhein, V., Kunc, H. P., & Naguib, M. (2004). Seasonal patterns of singing activity vary with time of day in the nightingale (*Luscinia megarhynchos*). *The Auk*, 121, 110–117.
- Amrhein, V., & Erne, N. (2006). Dawn singing reflects past territorial challenges in the winter wren. *Animal Behaviour*, 71, 1075–1080.
- Amrhein, V., Johannessen, L. E., Kristiansen, L., & Slagsvold, T. (2008). Reproductive strategy and singing activity: blue tit and great tit compared. *Behavioral Ecology and Sociobiology*, 62, 1633–1641.
- Ballentine, B., Badyaev, A., & Hill, G. E. (2003). Changes in song complexity correspond to periods of female fertility in blue grosbeaks (*Guiraca caerulea*). *Ethology*, 109, 55–66.
- Bédard, J., & LaPointe, G. (1984). Banding returns, arrival times, and site fidelity in the Savannah Sparrow. *The Wilson Bulletin*, 96, 196–205.
- Bruni, A., & Foote, J. R. (2014). Dawn singing of Eastern phoebes varies with breeding stage and brood number. *The Wilson Journal of Ornithology*, 126, 500–507.
- Burt, J. M., & Vehrencamp, S. L. (2005). Dawn chorus as an interactive communication network. *Animal Communication Networks*, 320–343.
- Catchpole, C. K. (1973). The Functions of Advertising Song in the Sedge Warbler (*Acrocephalus schoenobaenus*) and the Reed Warbler (*A. scirpaceus*). *Behaviour*, 46, 300–319.
- Cuthill, I. C., & Macdonald, W. A. (1990). Experimental manipulation of the dawn and dusk chorus in the blackbird *Turdus merula*. *Behavioral Ecology and Sociobiology*, 26, 209–216.
- Cuthill, I., & Hindmarsh, A. (1985). Increase in starling song activity with removal of mate. *Animal Behaviour*, 33, 326–328.
- Dalziel, A. H., & Cockburn, A. (2008). Dawn song in superb fairy-wrens: a bird that seeks extrapair copulations during the dawn chorus. *Animal Behaviour*, 75, 489–500.
- Demko, A. D., Reitsma, L. R., & Staicer, C. A. (2013). Two song categories in the Canada warbler (*Cardellina canadensis*). *The Auk*, 130, 609–616.
- Dixon, C. L. (1978). Breeding biology of the Savannah Sparrow on Kent Island. *The Auk*, 95, 235–246.

- Erne, N., & Amrhein, V. (2008). Long-term influence of simulated territorial intrusions on dawn and dusk singing in the Winter Wren: spring versus autumn. *Journal of Ornithology*, 149, 479–486.
- Foote, J. R., & Barber, C. a. (2009). Paired male Song Sparrows sing more when their mate is incubating. *The Wilson Journal of Ornithology*, 121, 819–822.
- Foote, J. R., Fitzsimmons, L. P., Mennill, D. J., & Ratcliffe, L. M. (2011). Male black-capped chickadees begin dawn chorusing earlier in response to simulated territorial insertions. *Animal Behaviour*, 81, 871–877.
- Gil, D., Graves, J. A., & Slater, P. J. B. (1999). Seasonal patterns of singing in the willow warbler: evidence against the fertility announcement hypothesis. *Animal Behaviour*, 58, 995–1000.
- Hanski, I., & Laurila, A. (1993). Variation in song rate during the breeding cycle of the chaffinch, *Fringilla coelebs*. *Ethology*, 93, 161–169.
- Johnson, L. S. & L. H. Kermott. (1991). The functions of song in male House Wrens (*Troglodytes aedon*). *Behaviour* 166:190-209.
- Kacelnik, A., & Krebs, J. R. (1983). The dawn chorus in the great tit (*Parus major*): proximate and ultimate causes. *Behaviour*, 83, 287–308.
- Kunc, H. P., Amrhein, V., & Naguib, M. (2005). Seasonal variation in dawn song characteristics in the common nightingale. *Animal Behaviour*, 70, 1265–1271.
- Lampe, H. M., & Espmark, Y. O. (1987). Singing activity and song pattern of the Redwing *Turdus iliacus* during the breeding season. *Ornis Scandinavica*, 18, 179–185.
- Leonard, M. (2008). An overview and comparative analysis of singing on the nest in North American birds. *Canadian Journal of Zoology*, 86, 1101–1110.
- Liu, W.-C. (2004). The effect of neighbours and females on dawn and daytime singing behaviours by male chipping sparrows. *Animal Behaviour*, 68, 39–44.
- Logan, C. A. (1983). Reproductively dependent song cyclicity in mated male mockingbirds (*Mimus polyglottos*). *The Auk*, 100, 404–413.
- Mace, R. (1987). The dawn chorus in the great tit *Parus major* is directly related to female fertility. *Nature*, 330, 745–746.
- Masco, C., Allesina, S., Mennill, D. J., & Pruett-Jones, S. (2016). The Song Overlap Null model Generator (SONG): a new tool for distinguishing between random and non-random song overlap. *Bioacoustics*, 25, 29–40.

- Moller, A. P. (1991). Why mated songbirds sing so much: mate guarding and male announcement of mate fertility status. *The American Naturalist*, 138, 994–1014.
- Murphy, M. T., Sexton, K., Dolan, A. C., & Redmond, L. J. (2008). Dawn song of the eastern kingbird: an honest signal of male quality? *Animal Behaviour*, 75, 1075–1084.
- Naguib, M., van Rooij, E. P., Snijders, L., & Van Oers, K. (2016). To sing or not to sing: seasonal changes in singing vary with personality in wild great tits. *Behavioral Ecology*, 27, 932–938.
- Ord, T. J. (2008). Dawn and dusk “chorus” in visually communicating Jamaican anole lizards. *The American Naturalist*, 172, 585–592.
- Pärt, T. (1991). Is dawn singing related to paternity insurance? The case of the collared flycatcher. *Animal Behaviour*, 41(3), 451–456.
- Poesel, A., Dabelsteen, T., Darden, S.-K., Delhey, K., & Peters, A. (2006). Territorial responses of male blue tits, *Cyanistes caeruleus*, to UV-manipulated neighbours. *Journal of Ornithology*, 148, 179–187.
- Potter, P. E. (1972). Territorial behavior in Savannah Sparrows in southeastern Michigan. *The Wilson Bulletin*, 48–59.
- Reid, M. L., & Weatherhead, P. J. (1990). Mate-choice criteria of Ipswich sparrows: the importance of variability. *Animal Behaviour*, 40, 538–544.
- Rodrigues, M. (1996). Song activity in the chiffchaff: territorial defence or mate guarding? *Animal Behaviour*, 51, 709–716.
- Ross Lein, M. (2007). Patterns of dawn singing by Buff-breasted Flycatchers. *Journal of Field Ornithology*, 78, 343–351.
- Slagsvold, T. (1977). Bird song activity in relation to breeding cycle, spring weather, and environmental phenology. *Ornis Scandinavica*, 8, 197–222.
- Slagsvold, T., Dale, S., & Sætre, G.-P. (1994). Dawn singing in the great tit (*Parus major*): mate attraction, mate guarding, or territorial defence? *Behaviour*, 131, 115–138.
- Staicer, C. A., Spector, D. A., & Horn, A. G. (1996). The dawn chorus and other diel patterns in acoustic signalling. In: Kroodsma D.E, Miller E.H, editors. *Ecology and evolution of acoustic communication in birds*. Cornell University, Laboratory of Ornithology, Ithaca, New York, USA, 426–453.
- Tufts, R. W. (1986). *Birds of Nova Scotia*. (3rd ed). Halifax, NS: Nimbus Publ. Ltd. and Nova Scotia Mus.

- Wheelwright, N., & Rising, J. (2008). Savannah Sparrow (*Passerculus sandwichensis*). Number 45. The Birds of North America. Cornell University, Laboratory of Ornithology, Ithaca, New York, USA.
- Wheelwright, N. T., & Mauck, R. A. (1998). Philopatry, natal dispersal, and inbreeding avoidance in an island population of Savannah Sparrows. *Ecology*, 79, 755–767.
- Wheelwright, N. T., Schultz, C. B., & Hodum, P. J. (1992). Polygyny and male parental care in Savannah sparrows: effects on female fitness. *Behavioral Ecology and Sociobiology*, 31, 279–289.
- Wheelwright, N. T., Swett, M. B., Levin, I. I., Kroodsma, D. E., Freeman-Gallant, C. R., & Williams, H. (2008). The influence of different tutor types on song learning in a natural bird population. *Animal Behaviour*, 75, 1479–1493.
- Williams, H., Levin, I. I., Norris, D. R., Newman, A. E. M., & Wheelwright, N. T. (2013). Three decades of cultural evolution in Savannah sparrow songs. *Animal Behaviour*, 85, 213–223.
- Woodworth, B. K., Mitchell, G. W., Norris, D. R., Francis, C. M., & Taylor, P. D. (2015). Patterns and correlates of songbird movements at an ecological barrier during autumn migration assessed using landscape-and regional-scale automated radiotelemetry. *Ibis*, 157, 326–339.
- Zhang, V. Y., Celis-Murillo, A., & Ward, M. P. (2016). Conveying information with one song type: changes in dawn song performance correspond to different female breeding stages. *Bioacoustics*, 25, 19–28.

Figures




Figure 2.1. Diel variation of the singing activity of Savannah Sparrows with respect to breeding stages. Breeding stages included: unpaired stage, fertile stage, incubation stage, hatchling stage, and fledgling stage ($n=34$ males). Note the peak in singing activity in the morning around 7:00 and absence of dusk chorus for the unpaired males, and the presence of dusk chorus in paired males (the dots represent means for each hour of the day). Bars show mean \pm SE.




Figure 2.2. Singing activity during the dawn chorus (30 min before sunrise) with respect to breeding stage (unpaired, fertile, incubation, hatchling and fledgling) (n=34). Unpaired males sang in the absence of females significantly more than any other stages. Bars show mean \pm SE, and levels that are not significantly different are represented with the same letter.




Figure 2.3. Singing activity during the daytime (time after sunrise and before sunset) with respect to five breeding stages (unpaired, fertile, incubation, hatchling and fledgling stages) (n=34). Unpaired sang significantly more than the fledgling stage. Bars show number \pm SE, and levels that are not significantly different are represented with the same letter.




Figure 2.4. Singing activity during the dusk chorus (time after sunset) with respect to five breeding stages (unpaired, fertile, incubation, hatchling and fledgling stages) ($n=34$). Males who were unpaired sang significantly less than males at all of the other stages, suggesting that dusk songs have a female-related function. Bars show mean \pm SE and levels that are not significantly different are represented with the same letter.

Supplementary Table 2.1. Results of diel singing activity LMM standard least square comparisons for all interactions.

	<i>t Ratio</i>	<i>P</i>
Treatment type		
Unpaired stage	8.95	<.0001*
Fertile stage	0.42	0.6765
Fledgling stage	-6.23	<.0001*
Hatchling stage	0.84	0.3993
Incubation stage	5.99	<.0001*
Hour		
Between hours	-4.75	<.0001*
Treatment type * hour		
Fertile stage*hour	6.38	<.0001*
Fledgling stage*hour	1.41	0.1584
Hatchling*hour	1.36	0.1737
Incubation*hour	3.29	0.0010*

Significant values shown with *

Supplementary Table 2.2. Results of seasonal singing activity LMM standard least square comparisons for all interactions.

	<i>t Ratio</i>	<i>P</i>
Treatment type		
Unpaired stage	5.09	<.0001*
Fertile stage	-2.37	0.0180*
Fledgling stage	0.93	0.3539
Hatchling stage	0.38	0.7018
Incubation stage	1.04	0.2988
Day		
Julian day	-3.30	0.0010*
Treatment type*Day		
Julian day*Fertile	1.66	0.0963
Julian day*Fledgling	-3.42	0.0006*
Julian day*Hatchling	-0.88	0.3803
Julian day*Incubation	0.07	0.9432

Significant values shown with *

CHAPTER 3: Quiet violence: Songbirds respond to playback simulated rivals using soft songs as aggressive signals of intention

Chapter summary

When animals compete over resources such as breeding territories, they often use signals to communicate their aggressive intentions. According to the hierarchical signalling hypothesis, animals progressively escalate interactions using specific signals to reveal their aggressive intentions towards conspecific rivals. In this study, we tested this hypothesis in male Savannah Sparrows, *Passerculus sandwichensis*, focusing on signals that precede physical attack against a territorial intruder. We simulated an intruder using song playback and a taxidermic model, and we determined which behaviors were associated with physical attack. Several previous studies have used this approach, and they provide evidence that many species produce songs of dramatically lower amplitude, or “soft songs”, as signals of aggressive intention. Savannah Sparrows, however, are not known to produce soft songs, and therefore they provide an interesting system for testing signals that predict attack. Out of 93 playback subjects, 23 males attacked the simulated intruder and 70 did not. To our surprise, Savannah Sparrows produced soft songs, a behaviour which had not previously been described for this well-studied songbird, and the number of soft songs was a significant predictor of attack on the simulated intruder. The number of “chip” calls also predicted attack on the simulated intruder, whereas three other measured behaviors (aggressive calls, wing waving and passes over the model) did not predict attack. Our study contributes to the growing body of research that supports the hierarchical signalling hypothesis (Searcy et

al., 2013; Searcy & Beecher, 2009) and corroborates with the idea that soft song is a widespread signal of aggression.

Introduction

Territorial interactions between rival animals routinely involve physical aggression. Given that aggressive encounters may involve costly physical attacks, or even death, many animals have evolved signals to facilitate communication during territorial interactions (Laidre & Johnstone, 2013; Vehrencamp, 2000). Some animals use graded signals to convey their intention to escalate or de-escalate encounters, including high-intensity signals of threat or low-intensity signals of submission (Searcy et al., 2013; Akçay et al., 2013; Hof & Podos, 2013). Although aggressive territorial signals have been documented in diverse animal taxa including mammals (August & Anderson, 1987; Behr et al., 2009; Clutton-Brock & Albon, 1979), birds (Searcy et al., 2006; Todt & Naguib, 2000), reptiles (Baird et al., 2013), frogs (Bee et al., 2000; Reichert & Gerhardt, 2013; Wagner, 1989), fish (Korzan & Fernald, 2007; Triefenbach & Zakon, 2008), mollusks (Schnell et al., 2016), and arthropods (Jonsson, Kravitz, & Heinrich, 2011; Tibbetts & Sheehan, 2011, Wagner, 1989), only recently have animal behaviourists come to understand the function of specific types of signals within a hierarchical signalling system of escalation and de-escalation. One particular experimental protocol helps to reveal the vocal and physical signals that predict attack; it involves the presentation of a simulated territorial rival (e.g. a taxidermic model, a mirror, or video playback) accompanied by acoustic stimuli (e.g. vocal playback; Searcy, et al., 2006). Under this experimental design, animals have the opportunity to attack the simulated rival, and we can gain insight into the function of specific signals by studying the behaviours that precede physical attack.

Hierarchical signalling models have been developed for animals across diverse taxa, including the signals associated with aggressive escalation in mammals (Poole, 1989), birds (Hof & Podos, 2013; Searcy & Beecher, 2009), fish (Triefenbach & Zakon, 2008), amphibians (Bortosky & Mathis, 2016), reptiles (Van Dyk & Evans, 2008), and arthropods (Schnell et al., 2016). For example, cuttlefish, *Sepia apama*, indicate their intention to attack a rival when they produce particular visual displays (specifically, shovel and lateral displays; Schnell et al., 2016), and knifefish, *Apteronotus leptorhynchus*, produce particular electrical signals as predictors of attack against rivals (specifically, with chirps and gradual frequency rises; Triefenbach & Zakon, 2008). Birds provide a model system for studying aggressive signals because they produce diverse acoustic signals, including elaborate and variable signals that are commonplace during territorial contests in many species (Todt & Naguib, 2000b). Acoustic signalling behaviours in birds include song overlapping (Helfer & Osiejuk, 2015; Naguib & Mennill, 2010), song matching (King & McGregor, 2016) variation in song rate (Baker et al., 2012), variation in trill rate and frequency bandwidth (Dubois et al., 2008), production of low-amplitude songs or soft songs (Ballentine et al., 2008; Grieves et al., 2015; Hof & Hazlett, 2010; Searcy et al., 2006; Xia et al., 2013), and production of non-song calls (Ballentine et al., 2008; Baker et al., 2012; Elie & Theunissen, 2015). A fine example of a model of hierarchical signalling comes from Little Blue Penguins, *Eudyptula minor* (Waas, 1991): Little Blue Penguins produce multiple graded signals (including growls and hisses), and when exposed to a taxidermic model of a conspecific rival, males produce growls to signal their intention to escalate an encounter, and hisses to signal their

intention to attack (Waas, 1991). The most well-studied system of territorial contest escalation in birds comes from Song Sparrows, *Melospiza melodia* (Akçay et al., 2013; Searcy & Beecher, 2009; Searcy et al., 2006). Male Song Sparrows escalate interactions by matching the songs of their rivals, or de-escalate interactions by switching to non-matching songs (Akçay et al., 2013). They produce low-amplitude songs, also known as “soft songs”, for their most-threatening signal, which precedes an attack on a rival (Searcy et al., 2006). Subsequent investigations in several bird taxa reveal that low-amplitude signals are often used as high-threat signals (reviewed in Searcy et al., 2014).

Low-amplitude signals are taxonomically widespread vocal behaviours found in birds (e.g. Dabelsteen et al., 1998; Reichard & Welklin, 2015), mammals (e.g. Gustison & Townsend, 2015), and invertebrates (e.g. Balenger, 2015). Historically overlooked in birds, soft vocalizations (soft songs and soft calls) are now known to be used by almost half of North American birds during territorial defence, courtship displays, and alarm, begging, or contact signalling (Reichard & Welklin, 2015). Thirty-one bird species are known to produce soft vocalizations as aggressive signals during displays of territorial defence in North America (Reichard & Welklin, 2015). Furthermore, males and females from six geographically widespread and distantly-related bird species produce soft songs (or soft calls) as a predictor of attack during territorial contests (Table 3.1), suggesting that these vocalizations are a taxonomically widespread, and possibly ancestral trait. Low-amplitude songs are interesting from a communication network perspective, because they may have evolved to limit the potential for eavesdropping; if low-amplitude songs are associated with the highest levels of aggression, animals may

benefit by minimizing the broadcast of information about their occurrence to nearby animals (Dabelsteen et al., 1998).

Songs are the main vocalization produced by songbirds during their daily activities on their breeding grounds, yet other vocalizations, such as non-song calls, are also produced during songbirds' daily activities. Calls, unlike songs, are simple and short non-learned vocalizations that serve various functions and can be gender-specific or used by both genders (Catchpole & Slater, 2008). In past studies that investigated vocal territorial strategies in birds, researchers mainly focused on song rate, song overlapping (a time-specific response to neighbouring songs), song-type matching (a pattern-specific response to neighbouring songs), song switching (a repertoire-related response in large song repertoire birds), or modulation of fine structural song elements as the primary aggressive signals in birds (Burt et al., 2001; DuBois et al., 2009; Naguib, 2005; Searcy et al. 2006). Surprisingly few studies have investigated calls as signalling strategies to escalate interactions (e.g. Ballentine et al., 2008), even though research on call signalling strategies can likely provide a deeper understanding of animal signalling behaviour.

In this study, we explore aggressive signalling during the territorial interactions in a temperate songbird, the Savannah Sparrow, *Passerculus sandwichensis*, by applying the experimental protocol developed by Searcy et al. (2006) involving playback and the presentation of a taxidermic model to test predictors of physical attack. We chose to study Savannah Sparrows using this approach because this species, in contrast to many other sparrows, is not known to produce soft songs (Searcy et al., 2006; Ballentine et al.,

2008; Wheelwright & Rising, 2008; Reichard & Anderson, 2015). Therefore, Savannah Sparrows provide an interesting system for testing predictors of physical attack. The goal of our study is to explore which signals are threatening signals that best predict physical attacks in Savannah Sparrows. We predicted that Savannah Sparrows would reliably indicate their intention to attack a rival during territorial interactions using singing behaviours, calling behaviours, or physical behaviours during aggressive interactions.

Material and methods

Study species and study site

Savannah Sparrows are migratory songbirds that live in open grasslands across North America (Wheelwright & Rising, 2008). We conducted our research at Bowdoin Scientific Station on Kent Island, New Brunswick, Canada (44°35'N 66°46'W), a 200-acre island in the Bay of Fundy. This island is home to a strongly philopatric population of Savannah Sparrows (Wheelwright & Mauck, 1998). The birds in this population have been studied for many decades, including careful annual studies over the last thirty years (Williams et al., 2013; Woodworth et al., 2017). On an annual basis, we capture and band birds inhabiting our study site, giving each animal a unique combination of coloured leg bands to facilitate recognition of individuals in the field. Over half of our playback subjects were individually-marked animals from our study site (n= 52 individuals). The remaining playback subjects were unbanded males from outside our study site (n=41 individuals). We distinguished between different unbanded individuals

on the basis of their territorial position and their individually distinctive song (as in Hick, Doucet, & Mennill, 2015; Moser-Purdy & Mennill, 2016).

Male Savannah Sparrows arrive from migration in mid-April (Woodworth et al., 2016) and begin to establish breeding territories using a variety of vocal signals (Potter, 1972). Males possess a single song type, which they are thought to learn within the first eight months of life (Wheelwright et al., 2008) and which remains constant throughout an individual's life (Williams et al., 2013). Males produce a variety of calls, including contact calls (the "chip" call), aggressive flight calls, as well as hostile notes (a.k.a. the "buzz" call; Gobeil, 1970; Wheelwright & Rising, 2008). In addition to their songs and calls, males also produce visual displays during territorial establishment including wing waving, flutter flights, tail raising, bill gaping, ptiloerection of crest feathers, parallel walking, and bouts of physical chases (Potter, 1972; Wheelwright & Rising, 2008). The relative importance of Savannah Sparrows' diverse vocalizations and physical displays remains poorly studied; these displays have the potential to comprise a graded system of aggressive signalling, although this has never been studied.

Playback experiment

We conducted our playback trials between mid-April and late-May 2016 between 6:00am and 12:00pm, when vocal interactions between territorial male Savannah Sparrows are commonplace. Prior to each playback trial, we positioned a taxidermic model of a Savannah Sparrow on top of a FoxPro Scorpion X1A loudspeaker, and we covered the model with a piece of camouflage-patterned cloth. We used four

different taxidermic models of similar size and appearance, perched in a normal posture, and we alternated between these models across trials. We placed small flags on either side of the model, at distances of 1m, 2m, and 4 m, to help an observer estimate the distance of playback subjects from the model.

A schematic representation of our experimental protocol is provided in Figure 3.1, showing two phases of playback: the initial period and the experimental period. Our playback experiment follows the general procedure of Searcy et al. (2006) involving the presentation of looped playback of male song and presentation of a taxidermic model. During the initial period, we attracted birds to the playback area by playing songs while the taxidermic model was hidden under a cloth. We played songs for 1 min during the initial period, followed by a period of 5 min of silent observation. At 5 minutes and 45 seconds, we removed the cloth from the taxidermic model by pulling a string. At 6 minutes, the experimental period began. During the experimental period, with the taxidermic model exposed, we played songs for 2 min followed by up to 20 min of silent observation, or until the subject attacked the taxidermic model. If the subject attacked the model, we considered the trial complete and we removed the model; this minimized stress to the playback subject and protected the models for later trials. We considered an attack any physical contact between the subject and the taxidermic model. Attacks were unambiguous: subjects frequently flew at the taxidermic model, often landing on top of the model and pecking at it. We aborted any trials where males in neighbouring territories also responded, and any trials that did not attract a subject during the initial five-minute period.

During the initial period, we played songs at an amplitude of 87 dB, and during the experimental period, we played songs at a quieter amplitude of 78 dB. We chose these two amplitudes to match those used by Searcy et al. (2006) in their study of Song Sparrows, a species related to Savannah Sparrows, and we decreased the amplitude to 78 dB like in Searcy et al. (2006) because soft vocalizations have been shown to act as aggressive signals during territorial contexts in other species (Table 3.1). We confirmed the amplitude of sounds from the loudspeaker in the field with a Casella CEL-240 sound level meter (Casella CEL Inc., Buffalo, NY, USA; C-weighting; fast setting), placed 1.0 m away from the speaker.

To record birds' responses to playback, we used two microphones connected to the two channels of a Marantz PMD661 digital recorder. We placed a Sennheiser ME66/K6 microphone on the ground approximately 45 cm from the speaker and taxidermic model, to capture any low-amplitude sounds produced by the subject during playback. We mounted a Sennheiser ME66/K6 microphone on a tripod 15-20 m away from the speaker and taxidermic model, to capture broadcast songs produced by the subject. An observer (I.G.M.) narrated the behaviours of the subject while they responded to the playback, by whispering into the second microphone while seated on the ground next to it.

Playback stimuli

We created playback stimuli from recordings we collected from male Savannah Sparrows in 2013-2015. We collected these recordings during spontaneous bouts of

singing by territorial birds, using a Marantz PMD661 digital recorder and a Sennheiser ME62/K6 microphone mounted in a Telinga parabola (sampling frequency: 44.1 kHz; sampling rate: 16 bit; WAVE format). Each playback subject received a different playback stimulus, and we chose stimuli that were recorded from a male at least 10 territories away from the subject ($\geq 700\text{m}$), ensuring that subjects were unfamiliar with the stimuli. From the original field recordings, we selected a single song with a high signal-to-noise ratio, and with little or no overlapping background sound. We then manipulated this song in Audition 3.0 software (Adobe, San Jose, CA, USA): we filtered out background noise with a high-pass filter of 1000 Hz and we normalized all sounds to the same amplitude (-1 dB). We then pasted each song into a longer sound file at a rate of six songs per minute, which is a typical song rate for Savannah Sparrows (based on natural song rate from focal recordings).

Analysis

We annotated our field recordings of the birds' playback responses in Syrinx-PC sound analysis software (Burt, Seattle, WA, USA), creating a second-by-second summary of the playback subjects' behaviour. We measured six aspects of each subject's behaviour: (1) number of broadcast songs, (2) number of soft songs, (3) number of "chip" calls, (4) number of aggressive calls (including both aggressive flight calls and buzzes (Wheelwright & Rising, 2008)), (5) number of times the bird waved its wings, and (6) number of passes over the taxidermic model. We included wing waving for comparison purposes with other playback studies of sparrows (Ballentine et al., 2008;

Searcy, Anderson, & Nowicki, 2006) even though our ability to maintain visual contact with subjects throughout the trials was constrained by the thick grassy habitat at the study site; our estimates of wing waving are likely underestimates, because subjects sometimes disappeared from our view for brief periods during playback. This also explains why we did not include other physical behaviours like bill gaping or ptiloerection into our analysis.

Early in our playback trials we found that subjects often produced low-amplitude “soft songs” (see Results); to differentiate between broadcast songs and soft songs, the field observer indicated whether each song was a “broadcast song” or “soft song” while describing the subjects’ behaviour. We confirmed whether each song was a broadcast or a soft song by comparing the recording from the two microphones used to record subjects’ responses; soft songs were typically visible in the recording from the directional microphone closest the taxidermic model, but faint or even absent from the directional microphone positioned next to the field observer. In several cases, we detected soft songs that were not detected by the recordist in the field, but that were identified during the process of annotation with Syrinx-PC, only appearing in the channel closest to the taxidermic model. In addition, soft songs differed from broadcast songs in that they had unique soft notes placed in between their regular notes. Those unique notes were not present in broadcast songs.

We focused our analysis on the aforementioned six variables during two time periods: (1) during the first 5 minutes of the initial period, and (2) during one minute

prior to the time that birds attacked the model during the experimental period. We were interested in comparing birds that attacked the taxidermic model versus birds that did not attack. To facilitate this comparison, we matched the timing of the one-minute-before-attack of birds that attacked the model to a corresponding one-minute period from randomly-chosen non-attackers, as in previous experiments that used this protocol (e.g. Searcy et al., 2006; Baker et al., 2012).

Statistical analysis

Our six response variables were not normally distributed, and could not be normalized through transformation due to a preponderance of zero values. We used a non-parametric Spearman's rank correlation to investigate the relationship between the six variables. We used non-parametric Mann-Whitney U tests to compare the behaviour of attackers versus non-attackers for each of the six response variables. In addition, we followed the approach used by Ballentine et al. (2008) to analyze similar playback-response data in Swamp Sparrows; we looked at the combinations of response variables that predicted attack, comparing attackers and non-attackers using a forward and backward stepwise discriminant function analysis.

Results

Of 93 male Savannah Sparrows that received playback accompanied by a taxidermic model simulating a conspecific rival, 23 males attacked the model and 70 did not. The playback subjects showed variation in their signalling behaviour, with most

subjects producing broadcast songs, chip calls, aggressive calls, and some subjects producing wing waving displays and passing over the taxidermic model. To our surprise, a substantial number of birds produced soft songs during the trials one minute before attack ($n=39$), a behaviour that had not previously been described in this species (Wheelwright & Rising, 2008; Reichard & Welkin, 2015). Soft songs were structurally similar to broadcast songs, but were composed of additional notes not present in broadcast songs. Although not measured in this study, broadcast songs were observed to reduce in number once the playback experiment started. Once the playback started, males tended to sing more soft songs and calls.

Initial period

In the initial period, we found no significant variation in the behaviour of birds that eventually attacked the model and those that did not (Fig. 3.2). Birds that eventually attacked the taxidermic model (hereafter “attackers”) and birds that did not (hereafter “non-attackers”) did not differ in number of broadcast songs ($U=0.22$, $P=0.83$), soft songs ($U=0.84$, $P=0.40$), chips ($U=0.87$, $P=0.38$), aggressive calls ($U=0.92$, $P=0.35$), bouts of wing waving ($U=1.0$, $P=0.32$), and passes over the speaker ($U=-0.20$, $P=0.84$). In the forward stepwise discriminant function analysis of the six response variables, aggressive calls were the only variable that discriminated between attackers and non-attackers (Wilk's $\lambda= 0.93$, $F_{1,90}= 6.92$, $P= 0.01$). A backward stepwise discriminant function analysis with the same six variables converged on identical results, with aggressive calls as the last variable to exit the model (Wilk's $\lambda= 0.93$, $F_{1,90}= 6.92$, $P= 0.01$)

One minute before attack

We compared behaviours between attackers and non-attackers by comparing the minute before Savannah Sparrows attacked the taxidermic model, to a random comparison minute for birds that did not attack. Attackers sang significantly more soft songs ($U=3.20$, $P=0.001$), and produced significantly more chip calls ($U=2.26$, $P=0.02$) than the non-attackers (Fig. 3.3). Attackers and non-attackers did not differ in the number of broadcast songs ($U=1.63$, $P=0.10$), aggressive calls ($U=0.26$, $P=0.79$), wing waves ($U=1.51$, $P=0.13$), or passes over the model ($U=1.3$, $P=0.19$) (Fig. 3.3). In the forward stepwise discriminant function analysis, amongst the six variables, soft songs entered the model first, followed by number of chips; this model discriminated between attackers and non-attackers (Wilk's $\lambda=0.92$, $F_{2,90}=3.90$, $P=0.02$). A backward stepwise discriminant function analysis converged on identical results, with soft songs and number of chip calls representing the last variables to exit the model (Wilk's $\lambda=0.92$, $F_{2,90}=3.90$, $P=0.02$).

Discussion

During simulated territorial intrusion, male Savannah Sparrows showed strong aggressive responses to simulated intruders. Twenty-five percent of males attacked a rival simulated with playback and a taxidermic model. Although soft songs have not previously been described in Savannah Sparrows, we found that territorial males regularly gave soft songs and that males that attacked sang significantly more soft songs prior to attack. In addition, attackers produced significantly more chip calls compared

with non-attackers prior to an attack. Consistent with one prediction of the hierarchical signalling hypothesis (Searcy et al., 2013; Searcy & Beecher, 2009), we conclude that soft songs and chip calls are signals that predict attack in Savannah Sparrows.

Beginning with a study of Song Sparrows just over a decade ago by Searcy et al. (2006), ten additional studies have used a playback-and-model-presentation design to explore signals that predict attack during territorial encounters in birds, including both Oscines and non-Oscines (Table 3.1). Collectively, they point towards the conclusion that soft songs reliably predict attack across diverse species. Deterring rivals with low amplitude vocalizations, instead of broadcast vocalizations, is an intriguing behaviour. Why do so many different species lower the amplitude of their songs during aggressive territorial interactions? One compelling explanation is the eavesdropping avoidance hypothesis (Dabelsteen et al., 1998) which proposes that individuals sing softly to avoid social eavesdropping by conspecific males or females, or interceptive eavesdropping by predators (McGregor, 1993; Searcy & Yasukawa, 2016). Eavesdroppers can impose high costs on signalers, both in the case of social eavesdropping (e.g. Mennill et al. 2003), and interceptive eavesdropping (e.g. Randall & Matocq, 1997), hence signalers develop strategies to avoid eavesdroppers (Dabelsteen, 2005; Dabelsteen et al., 1998). Two studies have tested the influence of predators on the amplitude of male songs (Akçay et al. 2016; Searcy & Nowicki, 2006), and both showed that under higher risk of predation, birds do not increase their soft song output. Although more evidence is needed, these results suggest that eavesdropping predators do not pressure birds to sing softly.

Similarly, one study tested whether eavesdropper males pressure fighting males to

lower their songs during dyadic interactions (Searcy & Nowicki, 2006). In this study, more males intruded on territories when soft songs were sung compared to when songs were broadcast, which contradicts the eavesdropping hypothesis. Indeed, when males hear a foreign song and hear no responses from their neighbors, males or floaters may visit the intruded territory to either obtain parts of the territory, or to chase the intruder away to reduce competition. Finally, the last hypothesis that could explain the lowering in amplitude of male songs is the use of eavesdropping by conspecific females (Mennill et al., 2003); but presently, no studies have yet investigated the role of females in the evolution of soft songs.

A second compelling explanation for why males produce soft songs during aggressive territorial interactions comes from the readiness hypothesis (Akçay et al, 2011). When intruders invade another male's territory, territorial males may benefit by following the intruder at a close distance to visually track them. When singing loudly, however, males often angle their head backward, and this may not allow for visual tracking. Therefore, males may be physically constrained to sing softly in the presence of a rival, to keep track of his whereabouts. Savannah Sparrows are known to have preferred singing perches (Wheelwright & Rising, 2008) on which they have been observed to raise their bill when broadcasting their songs (Clark, 1976). During our playback experiment, males that sang soft songs seemed to maintain a closed bill and did not tilt their head back while on the ground near the taxidermic model. These observations are consistent with the idea that visual tracking of territorial intruders may constrain males to signal softly.

If soft songs are aggressive signals that predict an imminent attack against a rival animal, how do animals maintain an honest signalling system, when a territorial animal could easily “bluff” by producing soft songs even in the absence of their willingness to escalate a territorial interaction? Historically, the honesty of vocal signals was a topic of controversy because aggressive vocal signals were thought to be easy-to-bluff due to their low production cost (Smith, 1974; Smith, 1979; Smith, 1982). In fact, vocal signals can honestly signal aggressive intentions if receivers retaliate against the animals producing aggressive signals (Anderson et al., 2012; Enquist, 1985; Vehrencamp, 2000) or if receivers recognize their bluffing rivals based on past interactions (Laidre, 2005; Van Rhijn & Vodegel, 1980). Under both scenarios, receivers ensure the honesty of aggressive signals by imposing a receiver-retaliation cost on the signallers (Anderson et al., 2012; Enquist, 1985; Vehrencamp, 2000). Future research can develop a deeper understanding of receiver-retaliation costs by adopting a receiver perspective (*sensu* Searcy & Beecher, 2009) by using playback simulating a territorial rival producing broadcast song versus soft song; with soft songs predicted to incite more aggressive reactions.

In addition to songs, calls play a central role during social interactions in birds (Gill et al. 2015). Many avian studies that explore acoustic signals during the breeding season focus on songs, but rarely explore calls. Nevertheless, similar to our study, calls were found to be a good predictor of attack in Black-capped Chickadees (Baker et al. 2012) but calls were not good predictor of attack in other studies like Swamp Sparrows (Ballentine et al 2008) and Black-throated Blue Warblers (Hof & Hazlett 2010). Chip calls

may be used in combination with soft songs as a signal of attack. However, because chip calls are also produced during other non-territorial interactions and are also produced by females (Wheelwright & Rising, 2008), chip calls may not be a reliable predictor of attack. Future research could investigate whether chip calls differ from soft songs in their functions by focusing on the rate of chip calls during territorial contexts.

In conclusion, our results show that Savannah Sparrows produce more low-amplitude soft songs before attacking a territorial rival, and produce more chip calls before attacking a rival. This pattern contributes to the growing body of work showing that many birds rely on low-amplitude aggressive signals to communicate their intention to escalate interactions. Future research should focus on testing the hypotheses that attempt to explain soft vocalizations, and should focus on elucidating the hierarchical signalling models of birds by investigating low-threat signals. In addition, other signals such as song rate, song frequency, song length, other song structural elements, and even behavioural or visual displays may act as predictors of attack and merit further investigations. Future studies should also focus on the structural differences between soft songs and broadcast songs, as some key elements of soft songs may indicate aggressiveness. Exploring the aggressive signals across a wide array of animals can expand our knowledge on animal communication and provide insight on how animals express aggression.

Literature cited

- Akçay, Ç., Clay, A., Campbell, S. E., & Beecher, M. D. (2016). The sparrow and the hawk: aggressive signalling under risk of predation. *Behavioral Ecology*, 27, 601–607.
- Akçay, Ç., Tom, M. E., Campbell, S. E., & Beecher, M. D. (2013). Song type matching is an honest early threat signal in a hierarchical animal communication system. *Proceedings of the Royal Society B: Biological Sciences*, 280, 20122517.
- Akçay, Ç., Tom, M. E., Holmes, D., Campbell, S. E., & Beecher, M. D. (2011). Sing softly and carry a big stick: Signals of aggressive intent in the song sparrow. *Animal Behaviour*, 82, 377–382.
- Anderson, R. C., Searcy, W. A., Hughes, M., & Nowicki, S. (2012). The receiver-dependent cost of soft song: A signal of aggressive intent in songbirds. *Animal Behaviour*, 83, 1443–1448.
- August, P. V., & Anderson, J. G. T. (1987). Mammal sounds and motivation-structural rules: a test of the hypothesis. *Journal of Mammalogy*, 68, 1–9.
- Baird, T. A., Hardy, I., & Briffa, M. (2013). Lizards and other reptiles as model systems for the study of contest behaviour. *Animal Contests*, 258–286.
- Balenger, S. L. (2015). Stridulated soft song by singing insects. *Animal Behaviour*, 105, 275–280.
- Ballentine, B., Searcy, W., & Nowicki, S. (2008). Reliable aggressive signalling in swamp sparrows. *Animal Behaviour*, 75, 693–703.
- Bee, M. a, Perrill, S. a, & Owen, P. C. (2000). Male green frogs lower the pitch of acoustic signals in defense of territories: a possible dishonest signal of size? *Behavioral Ecology*, 11, 169–177.
- Behr, O., Knörnschild, M., & Von Helversen, O. (2009). Territorial counter-singing in male sac-winged bats *Saccopteryx bilineata*: Low-frequency songs trigger a stronger response. *Behavioral Ecology and Sociobiology*, 63, 433–442.
- Bortosky, R., & Mathis, A. (2016). Honest signalling in aggressive contests between Ozark Zigzag Salamanders (*Plethodon angusticlavius*). *Copeia*, 2016, 60–66.
- Burt, J. M., Campbell, S. E., & Beecher, M. D. (2001). Song type matching as threat: a test using interactive playback. *Animal Behaviour*, 62, 1163–1170.
- Catchpole, C. K., & Slater, P. J. B. (2008). Bird song: Biological themes and variations.

- Clark, G. A. (1976). Taxonomic differences in head movements of vocalizing Passerines. *Bird-Banding*, 47, 123–128.
- Clutton-Brock, T., & Albon, S. (1979). The roaring of red deer and the evolution of honest advertisement. *Behaviour*, 69, 145–170.
- Dabelsteen, T. (2005). Public, private or anonymous? Facilitating and countering eavesdropping. *Animal Communication Networks*. Cambridge University Press, Cambridge, 38–62.
- Dabelsteen, T., McGregor, P. K., Lampe, H. M., Langmore, N. E., & Holland, J. (1998). Quiet song in Song birds: an overlooked phenomenon. *Bioacoustics*, 9, 89–105.
- DuBois, A. L., Nowicki, S., & Searcy, W. A. (2009). Swamp sparrows modulate vocal performance in an aggressive context. *Biology Letters*, 5, 163–165.
- Enquist, M. (1985). Communication during aggressive interactions with particular reference to variation in choice of behaviour. *Animal Behaviour*, 33, 1152–1161.
- Gobeil, R. E. (1970). Vocalizations of the Savannah Sparrow. *Bird-Banding*, 41, 18–21.
- Grieves, L. A., Logue, D. M., & Quinn, J. S. (2015). Ready to fight: Reliable predictors of attack in a cooperatively breeding, Non-Passerine bird. *Ethology*, 121, 1154–1165.
- Gustison, M. L., & Townsend, S. W. (2015). A survey of the context and structure of high- and low-amplitude calls in mammals. *Animal Behaviour*, 105, 281–288.
- Helfer, B., & Osiejuk, T. S. (2015). It takes all kinds in acoustic communication: A new perspective on the song overlapping phenomenon. *Ethology*, 121, 315–326.
- Hick, K. G., Doucet, S. M., & Mennill, D. J. (2015). Interspecific vocal discrimination in Neotropical wrens: responses to congeneric signals in sympatry and allopatry. *Animal Behaviour*, 109, 113–121.
- Hof, D., & Hazlett, N. (2010). Low-amplitude song predicts attack in a North American wood warbler. *Animal Behaviour*, 80, 821–828.
- Hof, D., & Podos, J. (2013). Escalation of aggressive vocal signals: a sequential playback study. *Proceedings of the Royal Society B*, 280 (August), 20131553.
- Jonsson, T., Kravitz, E. A., & Heinrich, R. (2011). Sound production during agonistic behavior of male *Drosophila melanogaster*. *Fly*, 5, 29–38.
- King, S. L., & McGregor, P. K. (2016). Vocal matching: the what, the why and the how. *Biology Letters*, 12, 20160666.

- Korzan, W. J., & Fernald, R. D. (2007). Territorial male color predicts agonistic behavior of conspecifics in a color polymorphic species. *Behavioral Ecology*, 18, 318–323.
- Laidre, M. (2005). Honest signalling of intentions: the importance of performance risk, receiver response, and prior knowledge of opponents in the threat displays of mandrills. *Behaviour*, 142, 455–476.
- Laidre, M., & Johnstone, R. (2013). Animal signals. *Current Biology*, 23, R829–R833.
- Maynard Smith, J. (1982). Do animals convey information about their intentions? *Journal of Theoretical Biology*, 97, 1–5.
- McGregor, P. K. (1993). Signalling in territorial systems: a context for individual identification, ranging and eavesdropping. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 340, 237–244.
- Mennill, D. J., Boag, P. T., & Ratcliffe, L. M. (2003). The reproductive choices of eavesdropping female black-capped chickadees, *Poecile atricapillus*. *Naturwissenschaften*, 90, 577–582.
- Moser-Purdy, C., & Mennill, D. J. (2016). Large vocal repertoires do not constrain the dear enemy effect: a playback experiment and comparative study of songbirds. *Animal Behaviour*, 118, 55–64.
- Naguib, M. (2005). Singing interactions in songbirds: implications for social relations and territorial settlement. *Animal Communication Networks*, 300–319.
- Naguib, M., & Mennill, D. J. (2010). The signal value of birdsong: empirical evidence suggests song overlapping is a signal. *Animal Behaviour*, 80, e11–e15.
- Poole, J. H. (1989). Announcing intent: the aggressive state of musth in African elephants. *Animal Behaviour*, 37, 153–155.
- Potter, P. E. (1972). Territorial behavior in Savannah Sparrows in southeastern Michigan. *The Wilson Bulletin*, 84, 48–59.
- Randall, J. A., & Matocq, M. D. (1997). Why do kangaroo rats (*Dipodomys spectabilis*) footdrum at snakes? *Behavioral Ecology*, 8, 404–413.
- Reichard, D. G., & Anderson, R. C. (2015). Why signal softly? The structure, function and evolutionary significance of low-amplitude signals. *Animal Behaviour*, 105, 253–265.
- Reichard, D. G., & Welklin, J. F. (2015). On the existence and potential functions of low-amplitude vocalizations in North American birds. *The Auk*, 132, 156–166.

- Reichert, M. S., & Gerhardt, H. C. (2013). Gray tree frogs, *Hyla versicolor*, give lower-frequency aggressive calls in more escalated contests. *Behavioral Ecology and Sociobiology*, 67, 795–804.
- Schnell, A. K., Smith, C. L., Hanlon, R. T., Hall, K. C., & Harcourt, R. (2016). Cuttlefish perform multiple agonistic displays to communicate a hierarchy of threats. *Behavioral Ecology and Sociobiology*, 70, 1643–1655.
- Searcy, W. A., Akçay, C., Nowicki, S., & Beecher, M. D. (2014). Aggressive signalling in Song Sparrows and other Songbirds. *Advances in the Study of Behavior*, 46, 89–125.
- Searcy, W. a., Anderson, R. C., & Nowicki, S. (2006). Bird song as a signal of aggressive intent. *Behavioral Ecology and Sociobiology*, 60, 234–241.
- Searcy, W. A., DuBois, A. L., Rivera-Cáceres, K., & Nowicki, S. (2013). A test of a hierarchical signalling model in song sparrows. *Animal Behaviour*, 86, 309–315.
- Searcy, W. A., & Nowicki, S. (2006). Signal interception and the use of soft song in aggressive interactions. *Ethology*, 112, 865–872.
- Searcy, W. A., & Yasukawa, K. (2016). Eavesdropping and cue denial in avian acoustic signals. *Animal Behaviour*, 124, 273–282.
- Searcy, W., Anderson, B., & Nowicki, S. (2006). Bird song as a signal of aggressive intent. *Behavioral Ecology and Sociobiology*, 60, 234–241.
- Searcy, W., & Beecher, M. D. (2009). Song as an aggressive signal in songbirds. *Animal Behaviour*, 78, 1281–1292.
- Tibbetts, E. A., & Sheehan, M. J. (2011). Facial Patterns are a Conventional Signal of Agonistic Ability in *Polistes exclamans* Paper Wasps. *Ethology*, 117, 1138–1146.
- Todt, D., & Naguib, M. (2000). Vocal Interactions in Birds: The Use of Song as a Model in Communication. *Advances in the Study of Behavior*, 29, 247–296.
- Triefenbach, F. A., & Zakon, H. H. (2008). Changes in signalling during agonistic interactions between male weakly electric knifefish, *Apteronotus leptorhynchus*. *Animal Behaviour*, 75, 1263–1272.
- Van Dyk, D. A., & Evans, C. S. (2008). Opponent assessment in lizards: Examining the effect of aggressive and submissive signals. *Behavioral Ecology*, 19, 895–901.
- Van Rhijn, J. G., & Vodegel, R. (1980). Being honest about one's intentions: an evolutionary stable strategy for animal conflicts. *Journal of Theoretical Biology*, 85, 623–641.

- Vehrencamp, S. L. (2000). Handicap, index, and conventional signal elements of bird song. *Animal Signals: Signalling and Signal Design in Animal Communication*, 277–300.
- Waas, J. R. (1991). Do little blue penguins signal their intentions during aggressive interactions with strangers? *Animal Behaviour*, 41, 375–382.
- Wagner, W. E. (1989). Graded aggressive signals in Blanchard's cricket frog: vocal responses to opponent proximity and size. *Animal Behaviour*, 38, 1025–1038.
- Wheelwright, N., & Rising, J. (2008). Savannah Sparrow (*Passerculus sandwichensis*). Number 45. *The Birds of North America*. Cornell University, Laboratory of Ornithology, Ithaca, New York, USA.
- Wheelwright, N. T., & Mauck, R. A. (1998). Philopatry, natal dispersal, and inbreeding avoidance in an island population of Savannah Sparrows. *Ecology*, 79, 755–767.
- Wheelwright, N. T., Swett, M. B., Levin, I. I., Kroodsma, D. E., Freeman-Gallant, C. R., & Williams, H. (2008). The influence of different tutor types on song learning in a natural bird population. *Animal Behaviour*, 75, 1479–1493.
- Williams, H., Levin, I. I., Norris, D. R., Newman, A. E. M., & Wheelwright, N. T. (2013). Three decades of cultural evolution in Savannah sparrow songs. *Animal Behaviour*, 85, 213–223.
- Woodworth, B. K., Mitchell, G. W., Norris, D. R., Francis, C. M., & Taylor, P. D. (2015). Patterns and correlates of songbird movements at an ecological barrier during autumn migration assessed using landscape-and regional-scale automated radiotelemetry. *Ibis*, 157, 326–339.
- Xia, C., Liu, J., Alström, P., Wu, Q., & Zhang, Y. (2013). Is the soft song of the brownish-flanked bush warbler an aggressive signal? *Ethology*, 119, 653–661.

Tables

Table 3.1. Summary of studies that experimentally investigated reliable predictors of attack in birds.

Measured behavioural responses ranged from soft vocalizations (soft songs or soft calls), visual displays (e.g. wing waving, throat inflation), aggressive calls (e.g. hoot), and increased signalling rate. Behavioural responses that were the most reliable predictor of attack are marked with “support”. If behavioural responses were not found to be the most reliable predictor of attack, they were marked with “no support”. Behavioural responses that were not tested, were marked as “not tested”. Studies that followed the playback methods or modified playback methods from Searcy et al. (2006) are indicated with an asterisk (*) and studies that used a taxidermic model are indicated with a dagger (†).

Species	Behavioural variables				Sample size	Number of attacks
	Soft vocalizations	Visual displays	Aggressive Calls	Signalling rate		
Little Blue Penguin (Waas 1991b) †	Not tested	Not tested	Support	Not tested	40	na
Black-capped Chickadee (Baker et al. 2012) * †	Not tested	Not tested	Support	Support	38	21
House Wren (Barnet et al. 2014) †	Not tested	Support	Not tested	Support	37	9
Smooth-billed Ani (Grievess & Logue 2015)†*	No support	Support	No support	Support	14	8
Song Sparrow (Searcy et al. 2006) * †	Support	No support	Not tested	Not tested	95	20
Swamp Sparrow (Ballentine et al. 2008)* †	Support	Support	No support	Not tested	31	9
Black-throated Blue Warbler (Hof & Hazlett 2010) * †	Support	Not tested	No support	No support	54	19
Corncrake (Ręk & Osiejuk 2011)	Support	Not tested	Not tested	No support	29	6
Brownish-Flanked Bush Warbler (Xia et al. 2013) *†	Support	Not tested	Not tested	Not tested	25	6
Savannah Sparrow (current study) * †	Support	No support	Support	Not tested	93	23

Figures




Figure 3.1. Diagram illustrating the playback experiment, based on the experimental design of Searcy et al. (2006). There are two periods: an initial period and an experimental period. The initial period consists of a playback phase (1min) followed by a silent phase (5min). The experimental period consists of a playback phase (2 min) followed by a silent phase that lasted up to 20 minutes or up to an attack with the taxidermic model. The taxidermic model was revealed at 5 minutes and 45 seconds. Vocalizations and behaviors were recorded from the time of the first song. We focused our analysis on two time periods: the initial period and the 1-minute prior to attack period.




Figure 3.2. Behaviors between attackers and non-attackers during the five-minute initial period of the playback experiment. The behaviours, shown here as the total number (mean \pm SE) did not differ statistically between attackers and non-attackers during this initial period, for (a) broadcast songs, (b) soft songs, (c) chips, (d) aggressive calls (including both aggressive flight calls and buzzes), (e) wing waving, and (f) passes over the taxidermic model.




Figure 3.3. Behaviors between attackers and non-attackers during the one-minute prior to attack period of the playback experiment during taxidermy model presentation trials. For non-attackers, the timing of the one minute before the attack for attackers was matched to a corresponding one-minute period from randomly-chosen non-attackers. Behaviours (mean number \pm SE) differed significantly between attackers and non-attackers with (b) soft songs, and (c) chips, but not with other behavioural measures, (a) broadcast songs, (d) aggressive calls (including both aggressive flight calls and buzzes), (e) wing waving, and (f) passes over the taxidermic model.

CHAPTER 4: General discussion

Bird songs are known to serve a dual function to both attract mates and defend territories (Catchpole & Slater, 2008), a pattern which is true across many other organisms across the animal kingdom (Bradbury & Vehrencamp, 1998). My thesis investigated those functions of bird song by focusing on variation in song output with time of the day, time of year, and breeding status (Chapter 2), and on the behaviour of animals during interactions with conspecific rivals (Chapter 3) in Savannah Sparrows, *Passerculus sandwichensis*.

In Chapter 2, I explored temporal variation in Savannah Sparrow song by analysing diel and seasonal variation in vocal activity. My analyses revealed that (1) male singing activity was highest in the month of May; (2) male singing activity was highest during the unpaired period and the incubation period; (3) territorial unpaired males had the strongest dawn chorus and morning singing activity, especially early in the year; and (4) territorial unpaired males did not sing, or sing very little, during the dusk chorus, but paired males sang significantly more during the dusk chorus during the fertile, incubation, hatchling, and fledgling periods. My results were consistent with other studies that show birds' singing activity varies with time of day and with time of year (Foote & Barber, 2009; Slagsvold, 1996; Topp & Mennill, 2008), and these patterns provide insight into the importance of heightened song output associated with breeding activities.

In Chapter 3, I investigated territorial signalling strategies in Savannah Sparrows by conducting a playback experiment that simulated territorial intrusion. Previous

studies on songbirds and non-songbirds showed that soft song serves as a signal that predicts attack in birds (e.g. Barbara Ballentine et al., 2003; Hof & Hazlett, 2010; Rek & Osiejuk, 2011; Searcy et al., 2006), although not all birds use this behaviour to express their intention to attack (e.g. Black-capped Chickadees, *Poecile atricapillus*, Baker et al., 2012). I focused on the signals that predict attack during aggressive contests in Savannah Sparrows by following the playback protocol as in Searcy et al., 2006. My analysis of behavioural responses showed that soft songs were a reliable predictor of attack during aggressive interactions between males, consistent with studies of other species of sparrow.

Chapter 2 of my thesis provides evidence that time of day, time of year, and breeding stage influences singing activity of birds and the function of avian vocal signals, highlighting the necessity of carefully considering the time of year and time of day when ornithologists collect data on bird behaviour. For example, the results for my playback study in Chapter 3 might have been different if conducted later in the breeding season when females were present, given the dramatic reduction in song output shown in Chapter 2. Furthermore, in Chapter 2, I showed that breeding status can affect singing activity and this should also be taken into consideration when assessing singing activity. For example, during the playback experiment I presented in Chapter 3, all males had the same breeding status (i.e. territorial unpaired males) and the observed behaviours may have changed once females arrived on the breeding grounds, especially if male propensity to produce soft songs varies with the presence of females. Consequently,

when conducting playback experiments, controlling for time of day, time of year, and breeding status will allow researchers to carefully control for variation in vocal output.

Investigations into the function of low-amplitude soft songs are relatively new and many questions about soft songs remain unanswered (Reichard & Welklin, 2015). To further understand the function of soft songs, future research could consider using seasonal variation studies. Just as the results of Chapter 2 of this thesis provided insight into the function of normal song, investigating temporal variation of soft song output could reveal interesting singing patterns that would provide insight into the function of soft songs. Similarly, analysis of diel and seasonal variation could be used to investigate aggressive call output variation. Just as our understanding of soft song is limited, the specific function of many calls and their importance in social interactions remain unclear and worthy of further investigation.

Additional future research could explore signalling strategies that facilitate escalation or de-escalation early in aggressive interactions, and provide a more comprehensive understanding of the hierarchical signalling strategies of Savannah Sparrows and other species. It is interesting to note that soft songs are used as predictor of attack in related sparrow species (e.g. Swamp Sparrows and Song Sparrows; Ballentine et al., 2008; Searcy et al., 2006) as well as in distantly related species (e.g. Smooth-billed Anis and Corn Crakes; Grieves et al., 2015; Rek & Osiejuk, 2011), suggesting that this signalling strategy may have evolved from a common ancestor but was lost in some lineages (e.g. Baker et al. 2012). Investigating the taxonomic diversity

of soft songs used during aggressive contexts would give further insight on how common this signalling strategy is used in birds.

Another question that remains answered is why soft songs evolved as reliable aggressive signals since they are so easily produced and easily cheatable? In order to answer this question, further future studies are needed to test several proposed hypotheses that explain why low amplitude songs are reliable aggressive signals, presumably through the cost of receiver retaliation (Akçay, et al. 2015).

In conclusion, my thesis provides strong evidence that the analysis of seasonal and diel variation in singing activity with respect to breeding stages is an important tool to understand how birds use their songs. My research showed that Savannah Sparrows have dynamic patterns of diel and seasonal vocal variation, where vocal output varies with the social context at different times of their breeding season. In addition, my research suggests that dawn choruses have a territorial function whereas dusk choruses may have a female-related function in this species. Furthermore, my thesis provides evidence that Savannah Sparrows use graded signalling strategies to interact during territorial interactions. Similar to past studies in other species, my research showed that Savannah Sparrows signal their intention to attack their rivals with soft songs. In addition to soft songs, Savannah Sparrows also produce chip calls that predicted attack. My research advances the field of animal communication and provides a basis for future research on signal function and on social interactions involving vocal signals.

Literature cited

- Akçay, Ç., Anderson, R. C., Nowicki, S., Beecher, M. D., & Searcy, W. A. (2015). Quiet threats : soft song as an aggressive signal in birds. *Animal Behaviour*, 105, 1–8.
- Baker, T. M., Wilson, D. R., & Mennill, D. J. (2012). Vocal signals predict attack during aggressive interactions in black-capped chickadees. *Animal Behaviour*, 84, 965–974.
- Ballentine, B., Badyaev, A., & Hill, G. E. (2003). Changes in song complexity correspond to periods of female fertility in blue grosbeaks (*Guiraca caerulea*). *Ethology*, 109, 55–66.
- Ballentine, B., Searcy, W., & Nowicki, S. (2008). Reliable aggressive signaling in swamp sparrows. *Animal Behaviour*, 75, 693–703.
- Bradbury, J. W., & Vehrencamp, S. L. (1998). *Principles of animal communication*. Sunderland, MA: Sinauer Associates.
- Catchpole, C., & Slater, P. J. B. (2008). *Bird song: Biological themes and variations*. Cambridge. England: Cambridge University Press.
- Foot, J. R., & Barber, C. a. (2009). Paired male Song Sparrows sing more when their mate is incubating. *The Wilson Journal of Ornithology*, 121, 819–822.
- Grievess, L. A., Logue, D. M., & Quinn, J. S. (2015). Ready to fight: Reliable predictors of attack in a cooperatively breeding, Non-Passerine bird. *Ethology*, 121, 1154–1165.
- Hof, D., & Hazlett, N. (2010). Low-amplitude song predicts attack in a North American wood warbler. *Animal Behaviour*, 80, 821–828.
- McGregor, P. K. (2005). *Animal communication networks*. Cambridge University Press.
- Rek, P., & Osiejuk, T. S. (2011). Nonpasserine bird produces soft calls and pays retaliation cost. *Behavioral Ecology*, 22, 657–662.
- Searcy, W. A., Anderson, R. C., & Nowicki, S. (2006). Bird song as a signal of aggressive intent. *Behavioral Ecology and Sociobiology*, 60, 234–241.
- Searcy, W. A., DuBois, A. L., Rivera-Cáceres, K., & Nowicki, S. (2013). A test of a hierarchical signalling model in song sparrows. *Animal Behaviour*, 86, 309–315.
- Searcy, W., Anderson, B., & Nowicki, S. (2006). Bird song as a signal of aggressive intent. *Behavioral Ecology and Sociobiology*, 60, 234–241.
- Searcy, W., & Beecher, M. D. (2009). Song as an aggressive signal in songbirds. *Animal Behaviour*, 78, 1281–1292.

Slagsvold, T. (1996). Dawn and dusk singing of male American Robins in relation to female behavior. *The Wilson Bulletin*, 108, 507–515.

Todt, D., & Naguib, M. (2000). Vocal interactions in birds: The use of song as a model in communication. *Advances in the Study of Behavior*, 29, 247-296.

Topp, S. M., & Mennill, D. J. (2008). Seasonal variation in the duetting behaviour of rufous-and-white wrens (*Thryothorus rufalbus*). *Behavioral Ecology and Sociobiology*, 62, 1107–1117.

Vita Auctoris

Name	Ines Moran
Place of Birth	Apt, Provence-Alpes-Côte d'Azur, FRANCE
Year of Birth	1989
Education	<p>University of Windsor, 2017 Windsor, Ontario Master's of Science, Biological Science</p> <p>University of British Columbia, 2015 Vancouver, British Columbia Bachelor of Science (with Distinctions), Major in Biology with Directed Studies in Biology</p> <p>Langara College, 2013 Vancouver, British Columbia Associate of Science</p>
Awards	<p>Animal Behavior Society Student Research Grant (2017)</p> <p>Department of Biology Chris Wysiekierski Memorial Scholarship (2017)</p> <p>Faculty of Science Impact Award (2017)</p> <p>American Ornithologists' Union Travel Award (2016)</p> <p>Nature Vancouver Award (2015)</p>
Conference presentations	<p>North American Ornithological Conference, Washington, USA, 2016</p> <p>Animal Behaviour Society Conference, Toronto, Canada, 2017</p> <p>American Ornithological Society / Society of Canadian Ornithologists Conference, Michigan, USA, 2017</p>